

**Example 124.** Consider the following system of (second-order) initial value problems:

$$\begin{aligned} y_1'' &= 2y_1' - 3y_2' + 7y_2 & y_1(0) &= 2, \quad y_1'(0) = 3, \quad y_2(0) = -1, \quad y_2'(0) = 1 \\ y_2'' &= 4y_1' + y_2' - 5y_1 \end{aligned}$$

Write it as a first-order initial value problem in the form  $\mathbf{y}' = M\mathbf{y}$ ,  $\mathbf{y}(0) = \mathbf{y}_0$ .

**Solution.** Introduce  $y_3 = y_1'$  and  $y_4 = y_2'$ . Then, the given system translates into

$$\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 7 & 2 & -3 \\ -5 & 0 & 4 & 1 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 2 \\ -1 \\ 3 \\ 1 \end{bmatrix}.$$

**Example 125.** Consider the following system of initial value problems:

$$\begin{aligned} y_1''' &= 2y_1'' - 3y_1' + 7y_2 & y_1(0) &= 2, \quad y_1'(0) = 3, \quad y_1''(0) = 4, \quad y_2(0) = -1, \quad y_2'(0) = 1 \\ y_2'' &= 4y_1' + y_2' + 5y_2 \end{aligned}$$

Write it as a first-order initial value problem in the form  $\mathbf{y}' = M\mathbf{y}$ ,  $\mathbf{y}(0) = \mathbf{y}_0$ .

**Solution.** Introduce  $y_3 = y_1'$ ,  $y_4 = y_1''$  and  $y_5 = y_2'$ . Then, the given system translates into

$$\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ -3 & 7 & 0 & 2 & 0 \\ 0 & 5 & 4 & 0 & 1 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 2 \\ -1 \\ 3 \\ 4 \\ 1 \end{bmatrix}.$$

### Solving systems of differential equations

**Example 126.** Determine the general solution to  $y_1' = 5y_1 + 4y_2 + e^{2x}$ ,  $y_2' = 8y_1 + y_2$ .

**Solution.** From the second equation it follows that  $y_1 = \frac{1}{8}(y_2' - y_2)$ . Using this in the first equation, we get  $\frac{1}{8}(y_2'' - y_2') = \frac{5}{8}(y_2' - y_2) + 4y_2 + e^{2x}$ . After multiplying with 8, this is  $y_2'' - y_2' = 5(y_2' - y_2) + 32y_2 + 8e^{2x}$ .

Simplified, this is  $y_2'' - 6y_2' - 27y_2 = 8e^{2x}$ , which is an inhomogeneous linear DE with constant coefficients which we know how to solve:

- Since the characteristic roots of the homogeneous DE are  $-3, 9$ , while the characteristic root for the inhomogeneous part is  $2$ , there must be a particular solution of the form  $y_p = Ce^{2x}$ . Plugging this  $y_p$  into the DE, we get  $y_p'' - 6y_p' - 27y_p = (4 - 6 \cdot 2 - 27)Ce^{2x} = -35Ce^{2x} \stackrel{!}{=} 8e^{2x}$ . Hence,  $C = -\frac{8}{35}$ .
- We therefore obtain  $y_2 = -\frac{8}{35}e^{2x} + C_1e^{-3x} + C_2e^{9x}$  as the general solution to the inhomogeneous DE.

We can then determine  $y_1$  as

$$\begin{aligned} y_1 &= \frac{1}{8}(y_2' - y_2) \\ &= \frac{1}{8} \left( -\frac{16}{35}e^{2x} - 3C_1e^{-3x} + 9C_2e^{9x} + \frac{8}{35}e^{2x} - C_1e^{-3x} - C_2e^{9x} \right) \\ &= -\frac{1}{35}e^{2x} - \frac{1}{2}C_1e^{-3x} + C_2e^{9x}. \end{aligned}$$

**Solution. (alternative)** We can also start with  $y_2 = \frac{1}{4}y_1' - \frac{5}{4}y_1 - \frac{1}{4}e^{2x}$  (from the first equation), although the algebra will require a little more work. In that case, we have  $y_2' = \frac{1}{4}y_1'' - \frac{5}{4}y_1' - \frac{1}{2}e^{2x}$ . Using this in the second equation, we get  $\frac{1}{4}y_1'' - \frac{5}{4}y_1' - \frac{1}{2}e^{2x} = 8y_1 + \frac{1}{4}y_1' - \frac{5}{4}y_1 - \frac{1}{4}e^{2x}$ .

Simplified, this is  $y_1'' - 6y_1' - 27y_1 = e^{2x}$ , which is an inhomogeneous linear DE with constant coefficients which we know how to solve:

- Since the characteristic roots of the homogeneous DE are  $-3, 9$ , while the characteristic root for the inhomogeneous part is  $2$ , there must be a particular solution of the form  $y_p = Ce^{2x}$ . Plugging this  $y_p$  into the DE, we get  $y_p'' - 6y_p' - 27y_p = (4 - 6 \cdot 2 - 27)Ce^{2x} = -35Ce^{2x} \stackrel{!}{=} e^{2x}$ . Hence,  $C = -\frac{1}{35}$ .
- We therefore obtain  $y_1 = -\frac{1}{35}e^{2x} + C_1e^{-3x} + C_2e^{9x}$  as the general solution to the inhomogeneous DE.

We can then determine  $y_2$  as

$$\begin{aligned} y_2 &= \frac{1}{4}y_1' - \frac{5}{4}y_1 - \frac{1}{4}e^{2x} \\ &= \frac{1}{4}\left(-\frac{2}{35}e^{2x} - 3C_1e^{-3x} + 9C_2e^{9x}\right) - \frac{5}{4}\left(-\frac{1}{35}e^{2x} + C_1e^{-3x} + C_2e^{9x}\right) - \frac{1}{4}e^{2x} \\ &= -\frac{8}{35}e^{2x} - 2C_1e^{-3x} + C_2e^{9x}. \end{aligned}$$

**Important.** Make sure you can explain why both of our solutions are equivalent!

### Example 127.

- (a) Determine the general solution to  $y_1' = 5y_1 + 4y_2$ ,  $y_2' = 8y_1 + y_2$ .

**Comment.** In matrix form, with  $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ , this is  $\mathbf{y}' = \begin{bmatrix} 5 & 4 \\ 8 & 1 \end{bmatrix} \mathbf{y}$ .

- (b) Solve the IVP  $y_1' = 5y_1 + 4y_2$ ,  $y_2' = 8y_1 + y_2$ ,  $y_1(0) = 0$ ,  $y_2(0) = 1$ .

**Solution.**

- (a) Note that this is the homogeneous system corresponding to the previous problem. It therefore follows from our previous solution (the latter one) that  $y_1 = C_1e^{-3x} + C_2e^{9x}$  and  $y_2 = -2C_1e^{-3x} + C_2e^{9x}$  is the general solution of the homogeneous system.
- (b) We already have the general solutions  $y_1, y_2$  to the two DEs. We need to determine the (unique) values of  $C_1$  and  $C_2$  to match the initial conditions:  $y_1(0) = C_1 + C_2 \stackrel{!}{=} 0$ ,  $y_2(0) = -2C_1 + C_2 \stackrel{!}{=} 1$ . We solve these two equations and find  $C_1 = -\frac{1}{3}$  and  $C_2 = \frac{1}{3}$ . The unique solution to the IVP therefore is  $y_1 = -\frac{1}{3}e^{-3x} + \frac{1}{3}e^{9x}$  and  $y_2 = \frac{2}{3}e^{-3x} + \frac{1}{3}e^{9x}$ .