

We now revisit and finish Example 120:

Example 128. Consider two brine tanks. Initially, tank T_1 is filled with 24gal water containing 3lb salt, and tank T_2 with 9gal pure water.

- T_1 is being filled with 54gal/min water containing 0.5lb/gal salt.
- 72gal/min well-mixed solution flows out of T_1 into T_2 .
- 18gal/min well-mixed solution flows out of T_2 into T_1 .
- Finally, 54gal/min well-mixed solution is leaving T_2 .

How much salt is in the tanks after t minutes?

Solution. Note that the amount of water in each tank is constant because the flows balance each other.

Let $y_i(t)$ denote the amount of salt (in lb) in tank T_i after time t (in min). In the time interval $[t, t + \Delta t]$:

$$\Delta y_1 \approx 54 \cdot \frac{1}{2} \cdot \Delta t - 72 \cdot \frac{y_1}{24} \cdot \Delta t + 18 \cdot \frac{y_2}{9} \cdot \Delta t, \text{ so } y_1' = 27 - 3y_1 + 2y_2. \text{ Also, } y_1(0) = 3.$$

$$\Delta y_2 \approx 72 \cdot \frac{y_1}{24} \cdot \Delta t - (18 + 54) \cdot \frac{y_2}{9} \cdot \Delta t, \text{ so } y_2' = 3y_1 - 8y_2. \text{ Also, } y_2(0) = 0.$$

In conclusion, we have obtained the system of equations

$$\begin{aligned} y_1' &= -3y_1 + 2y_2 + 27, & y_1(0) &= 3, \\ y_2' &= 3y_1 - 8y_2, & y_2(0) &= 0. \end{aligned}$$

One strategy to solve this system is to first combine the two DEs to get a single equation for y_1 .

- From the first DE, we get $y_2 = \frac{1}{2}y_1' + \frac{3}{2}y_1 - \frac{27}{2}$.
- Using this in the second DE, we obtain $\left(\frac{1}{2}y_1' + \frac{3}{2}y_1 - \frac{27}{2}\right)' = 3y_1 - 8\left(\frac{1}{2}y_1' + \frac{3}{2}y_1 - \frac{27}{2}\right)$.
Simplified, this is $y_1'' + 11y_1' + 18y_1 = 216$.
- We already have the initial condition $y_1(0) = 3$. We get a second one by combining $y_2 = \frac{1}{2}y_1' + \frac{3}{2}y_1 - \frac{27}{2}$ with $y_2(0) = 0$ to get $0 = y_2(0) = \frac{1}{2}y_1'(0) + \frac{3}{2}y_1(0) - \frac{27}{2} = \frac{1}{2}y_1'(0) - 9$, which simplifies to $y_1'(0) = 18$.
- The IVP $y_1'' + 11y_1' + 18y_1 = 216$ with initial conditions $y_1(0) = 3$ and $y_1'(0) = 18$ is one that we can solve!
 - The general solution of the corresponding homogeneous equation is $y_h = C_1e^{-2t} + C_2e^{-9t}$.
 - The simplest particular solution is of the form $y_p = C$. Plugging into the DE, we find $y_p = \frac{216}{18} = 12$.
 - Hence, the general solution to the (inhomogeneous) DE is $y_1(x) = 12 + C_1e^{-2t} + C_2e^{-9t}$.
We then use the initial conditions $y_1(0) = 12 + C_1 + C_2 \stackrel{!}{=} 3$, $y_1'(0) = -2C_1 - 9C_2 \stackrel{!}{=} 18$ to find that for the unique solution of the IVP $C_1 = -9$, $C_2 = 0$.

The unique solution is $y_1(t) = 12 - 9e^{-2t}$.

- It follows that $y_2 = \frac{1}{2}y_1' + \frac{3}{2}y_1 - \frac{27}{2} = \frac{9}{2} - \frac{9}{2}e^{-2t}$.

Note. We could have found a particular solution with less calculations by observing (looking at the characteristic roots of the homogeneous DE and the inhomogeneous part) that there must be a solution of the form $\mathbf{y}_p(t) = \mathbf{a}$. We can then find \mathbf{a} by plugging into the differential equation. However, noticing that, for a constant solution, each tank has to have a constant concentration of 0.5lb/gal of salt, we find $\mathbf{y}_p(t) = \begin{bmatrix} 12 \\ 4.5 \end{bmatrix}$.

Example 129. (extra) Three brine tanks T_1, T_2, T_3 .

T_1 contains 20gal water with 10lb salt, T_2 40gal pure water, T_3 50gal water with 30lb salt.

T_1 is filled with 10gal/min water with 2lb/gal salt. 10gal/min well-mixed solution flows out of T_1 into T_2 . Also, 10gal/min well-mixed solution flows out of T_2 into T_3 . Finally, 10gal/min well-mixed solution is leaving T_3 . How much salt is in the tanks after t minutes?

Solution. Let $y_i(t)$ denote the amount of salt (in lb) in tank T_i after time t (in min).

In the time interval $[t, t + \Delta t]$:

$$\Delta y_1 \approx 10 \cdot 2 \cdot \Delta t - 10 \frac{y_1}{20} \cdot \Delta t, \text{ so } y_1' = 20 - \frac{1}{2}y_1. \text{ Also, } y_1(0) = 10.$$

$$\Delta y_2 \approx 10 \cdot \frac{y_1}{20} \cdot \Delta t - 10 \frac{y_2}{40} \cdot \Delta t, \text{ so } y_2' = \frac{1}{2}y_1 - \frac{1}{4}y_2. \text{ Also, } y_2(0) = 0.$$

$$\Delta y_3 \approx 10 \cdot \frac{y_2}{40} \cdot \Delta t - 10 \frac{y_3}{50} \cdot \Delta t, \text{ so } y_3' = \frac{1}{4}y_2 - \frac{1}{5}y_3. \text{ Also, } y_3(0) = 30.$$

Using matrix notation and writing $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$, this is $\mathbf{y}' = \begin{bmatrix} -1/2 & 0 & 0 \\ 1/2 & -1/4 & 0 \\ 0 & 1/4 & -1/5 \end{bmatrix} \mathbf{y} + \begin{bmatrix} 20 \\ 0 \\ 0 \end{bmatrix}$, $\mathbf{y}(0) = \begin{bmatrix} 10 \\ 0 \\ 30 \end{bmatrix}$.

We can actually solve this IVP!

[Do it! Start by finding y_1 from the first DE, then move on to $y_2 \dots$]

Here, we content ourselves with finding a particular solution (and ignoring the initial conditions). The method of undetermined coefficients tells us that there is a solution of the form $\mathbf{y}_p(t) = \mathbf{a}$. Of course, we can find \mathbf{a} by plugging into the differential equation. However, noticing that, for a constant solution, each tank has to have a concentration of 2lb/gal of salt, we find $\mathbf{y}_p = (40, 80, 100)$ without calculation.

Example 130. (April Fools!) Let $a = b$. Then $a^2 = ab$, so $a^2 + a^2 = a^2 + ab$ or $2a^2 = a^2 + ab$. Hence, $2a^2 - 2ab = a^2 - ab$ or $2(a^2 - ab) = a^2 - ab$. Cancelling, we arrive at $2 = 1$.

[Can you see the foul but disguised division by zero?!]