

A crash course in computing determinants

Review. The **determinant** of A , written as $\det(A)$ or $|A|$, is a number with the property that:

$$\begin{aligned} \det(A) \neq 0 &\iff A \text{ is invertible} \\ &\iff Ax = b \text{ has a (unique) solution } x \text{ (for all } b) \\ &\iff Ax = 0 \text{ is only solved by } x = 0 \end{aligned}$$

Example 41. $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$

Example 42. Compute $\begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix}$ by **cofactor expansion**.

Solution. We expand by the first row:

$$\begin{aligned} \begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} &= 1 \cdot \begin{vmatrix} + & & \\ & -1 & 2 \\ & 0 & 1 \end{vmatrix} - 2 \cdot \begin{vmatrix} & - & \\ 3 & & 2 \\ 2 & & 1 \end{vmatrix} + 0 \cdot \begin{vmatrix} & & + \\ 3 & -1 & \\ 2 & 0 & \end{vmatrix} \\ &\stackrel{\text{i.e.}}{=} 1 \cdot \begin{vmatrix} -1 & 2 \\ 0 & 1 \end{vmatrix} - 2 \cdot \begin{vmatrix} 3 & 2 \\ 2 & 1 \end{vmatrix} + 0 \cdot \begin{vmatrix} 3 & -1 \\ 2 & 0 \end{vmatrix} = 1 \cdot (-1) - 2 \cdot (-1) + 0 = 1 \end{aligned}$$

Each term in the cofactor expansion is ± 1 times an entry times a smaller determinant (row and column of entry deleted).

The ± 1 is assigned to each entry according to $\begin{bmatrix} + & - & + & \dots \\ - & + & - & \\ + & - & + & \\ \vdots & & & \ddots \end{bmatrix}$.

Solution. We expand by the second column:

$$\begin{aligned} \begin{vmatrix} 1 & 2 & 0 \\ 3 & -1 & 2 \\ 2 & 0 & 1 \end{vmatrix} &= -2 \cdot \begin{vmatrix} & - & \\ 3 & & 2 \\ 2 & & 1 \end{vmatrix} + (-1) \cdot \begin{vmatrix} 1 & & 0 \\ & + & \\ 2 & & 1 \end{vmatrix} - 0 \cdot \begin{vmatrix} 1 & & 0 \\ 3 & & 2 \\ & - & \end{vmatrix} \\ &= -2 \cdot (-1) + (-1) \cdot 1 - 0 = 1 \end{aligned}$$

Example 43. Compute $\begin{vmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 2 & 0 & 8 & 5 \end{vmatrix}$.

Solution. We can expand by the second column:

$$\begin{vmatrix} 1 & 0 & 3 & 4 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 2 & 1 \\ 2 & 0 & 8 & 5 \end{vmatrix} = -0 \begin{vmatrix} 0 & 1 & 5 \\ 0 & 2 & 1 \\ 2 & 8 & 5 \end{vmatrix} + 2 \begin{vmatrix} 1 & 3 & 4 \\ 0 & 2 & 1 \\ 2 & 8 & 5 \end{vmatrix} - 0 \begin{vmatrix} 1 & 3 & 4 \\ 0 & 1 & 5 \\ 2 & 8 & 5 \end{vmatrix} + 0 \begin{vmatrix} 1 & 3 & 4 \\ 0 & 1 & 5 \\ 0 & 2 & 1 \end{vmatrix}$$

[Of course, you don't have to spell out the 3×3 matrices that get multiplied with 0.]

We can compute the remaining 3×3 matrix in any way we prefer. One option is to expand by the first column:

$$2 \begin{vmatrix} 1 & 3 & 4 \\ 0 & 2 & 1 \\ 2 & 8 & 5 \end{vmatrix} = 2 \left(+1 \begin{vmatrix} 2 & 1 \\ 8 & 5 \end{vmatrix} + 2 \begin{vmatrix} 3 & 4 \\ 2 & 1 \end{vmatrix} \right) = 2(1 \cdot 2 + 2 \cdot (-5)) = -16$$

Comment. For cofactor expansion, choosing to expand by the second column is the best choice because this column has more zeros than any other column or row.

Solving linear recurrences with constant coefficients

Motivation: Fibonacci numbers

The numbers $0, 1, 1, 2, 3, 5, 8, 13, 21, 34, \dots$ are called **Fibonacci numbers**.

They are defined by the recursion $F_{n+1} = F_n + F_{n-1}$ and $F_0 = 0, F_1 = 1$.

How fast are they growing?

Have a look at ratios of Fibonacci numbers: $\frac{2}{1} = 2, \frac{3}{2} = 1.5, \frac{5}{3} \approx 1.667, \frac{8}{5} = 1.6, \frac{13}{8} = 1.625, \frac{21}{13} = 1.615, \frac{34}{21} = 1.619, \dots$

These ratios approach the **golden ratio** $\varphi = \frac{1+\sqrt{5}}{2} = 1.618\dots$

In other words, it appears that $\lim_{n \rightarrow \infty} \frac{F_{n+1}}{F_n} = \frac{1+\sqrt{5}}{2}$.

We will soon understand where this is coming from.

We can derive all of that using the same ideas as in the case of linear differential equations. The crucial observation that we can write the recursion in operator form:

$$F_{n+1} = F_n + F_{n-1} \quad \text{is equivalent to} \quad (N^2 - N - 1)F_n = 0.$$

Here, N is the shift operator: $Na_n = a_{n+1}$.

Comment. Recurrence equations are discrete analogs of differential equations.

For instance, recall that $f'(x) \approx f(x+1) - f(x)$ so that D is approximated by $N - 1$.

Example 44. Find the general solution to the recursion $a_{n+1} = 7a_n$.

Solution. Note that $a_n = 7a_{n-1} = 7 \cdot 7a_{n-2} = \dots = 7^n a_0$.

Hence, the general solution is $a_n = C \cdot 7^n$.

Comment. This is analogous to $y' = 7y$ having the general solution $y(x) = Ce^{7x}$.