

(reflections) Suppose that A is the matrix for reflecting through the plane W in 3-space.

- The 1-eigenspace of A is W . (dimension 2)
- The -1 -eigenspace of A is W^\perp . (dimension 1)

Why? By definition, the 1-eigenspace of A consists of those vectors that get reflected to themselves. But those are precisely the vectors in the plane W (only vectors on the plane are unchanged by the reflection). On the other hand, the -1 -eigenspace consists of those vectors v that get reflected to $-v$ (the exact opposite direction). These are precisely the vectors orthogonal to the plane.

Comment. In this context, the line W^\perp is often called the **normal line** of the plane W .

Example 108. Let A be the matrix for reflecting through the plane $W = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}\right\}$.

- (a) Diagonalize A (without first computing A) as $A = PDP^T$.
- (b) Is A invertible, orthogonal, symmetric?

Solution.

- (a) The eigenvalues of A are $1, 1, -1$. The 1-eigenspace of A is W , and the -1 -eigenspace is W^\perp .

In order to achieve a diagonalization PDP^T we need to choose P to be orthogonal (which we can do here because the eigenspaces are orthogonal).

As in the previous example, $W^\perp = \text{span}\left\{\begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}\right\}$.

We therefore choose $D = \begin{bmatrix} 1 & & \\ & 1 & \\ & & -1 \end{bmatrix}$ and, after normalizing columns, $P = \begin{bmatrix} 1/\sqrt{3} & -1/\sqrt{2} & 1/\sqrt{6} \\ 1/\sqrt{3} & 0 & -2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \end{bmatrix}$.

- (b) A is invertible (because 0 is not an eigenvalue).

By the same reasoning as in the previous example, A is symmetric.

Finally, note that $A^2 = I$ (reflecting twice isn't doing anything), so that $A^{-1} = A$. It follows that A is orthogonal, because $A^{-1} = A = A^T$.

By the way. Multiplying out $A = PDP^T$, we can find that $A = \frac{1}{3} \begin{bmatrix} 2 & 2 & -1 \\ 2 & -1 & 2 \\ -1 & 2 & 2 \end{bmatrix}$.

Comment. Similarly, a $n \times n$ matrix corresponds to a reflection (through a hyperplane) if and only if it has a $(n-1)$ -dimensional 1-eigenspace and a 1-dimensional -1 -eigenspace and these two spaces are orthogonal.

An alternative way of computing reflection matrices. Realize that, if n is the vector orthogonal to the plane (i.e. n is the normal vector of the plane), then reflecting v means sending it to $v - 2(\text{projection of } v \text{ onto } n)$.

We already observed that $n = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$.

Hence, the reflection of v is $v - 2(\text{projection of } v \text{ onto } n) = v - 2n \frac{n \cdot v}{n \cdot n} = v - 2 \frac{nn^T v}{n^T n} = \left(I - 2 \frac{nn^T}{n^T n}\right)v$.

Accordingly, the reflection matrix is $A = I - 2 \frac{nn^T}{n^T n} = \begin{bmatrix} 1 & & \\ & 1 & \\ & & 1 \end{bmatrix} - \frac{2}{6} \begin{bmatrix} 1 & -2 & 1 \\ -2 & 4 & -2 \\ 1 & -2 & 1 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 2 & 2 & -1 \\ 2 & -1 & 2 \\ -1 & 2 & 2 \end{bmatrix}$.

Comment. In other words, we got A from subtracting 2 times the projection matrix onto n from I .

Application: Linear differential equations

Example 109. (warmup) Solve the differential equation (DE) $y' = 2$.

Solution. From calculus, we know that the solutions are of the form $y(t) = 2t + C$.

Comment. To get a unique solution, we need to specify additional information, like an initial condition.

Example 110. (warmup) Solve the initial value problem (IVP) $y' = 2$, $y(0) = 1$.

Solution. This has the unique solution $y(t) = 2t + 1$.

Example 111. Which functions $y(t)$ satisfy the differential equation $y' = y$?

Solution. $y(t) = e^t$ and, more generally, $y(t) = Ce^t$. (And nothing else.)

(exponential function) e^t is the unique solution to $y' = y$, $y(0) = 1$.

From here, it follows that $e^t = 1 + t + \frac{t^2}{2!} + \frac{t^3}{3!} + \dots$

The latter is the Taylor series for e^t at $t = 0$ that we have seen in Calculus II.

Important note. We can actually construct this infinite sum directly from $y' = y$ and $y(0) = 1$.

Indeed, observe how each term, when differentiated, produces the term before it. For instance, $\frac{d}{dt} \frac{t^3}{3!} = \frac{t^2}{2!}$.

Example 112. Show that the differential equation $y' = 3y$ is solved by $y(t) = Ce^{3t}$.

Solution. Indeed, if $y(t) = Ce^{3t}$, then $y'(t) = 3Ce^{3t} = 3y(t)$.

Comment. It is important to realize that we can always easily check whether a function solves a differential equation. This means that (although you might be unfamiliar with the techniques for solving) you can use computer algebra systems like Sage to solve differential equations without trust issues.

Example 113. Solve the differential equation $y' = ay$ with initial condition $y(0) = y_0$.

Solution. As in the previous example, the general solution to $y' = ay$ is $y(t) = Ce^{at}$.

Since $y(0) = Ce^0 = C = y_0$, we conclude that the unique solution to the IVP is $y(t) = e^{at}y_0$.

Comment. It looks silly to write $e^{at}y_0$ instead of y_0e^{at} here, but we will soon replace the number a with a matrix A , and in that case only $e^{At}y_0$ makes sense.

Example 114. Our goal is to solve (systems of) differential equations like:

$$\begin{aligned}y_1' &= 2y_1 & y_1(0) &= 1 \\y_2' &= -y_1 + 3y_2 + y_3 & y_2(0) &= 0 \\y_3' &= -y_1 + y_2 + 3y_3 & y_3(0) &= 2\end{aligned}$$

In matrix form, this becomes

$$\mathbf{y}' = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}.$$

The key idea will be to solve $\mathbf{y}' = A\mathbf{y}$ by introducing e^{At} .

Theorem 115. The solution to $\mathbf{y}' = A\mathbf{y}$, $\mathbf{y}(0) = \mathbf{y}_0$ is $\mathbf{y}(t) = e^{At}\mathbf{y}_0$.

Recall from Example 113 that the solution to $y' = ay$, $y(0) = y_0$ is $y(t) = e^{at}y_0$. Here, however, At is a matrix and so we need to make sense of the matrix exponential. Next time, we will define e^A by the familiar Taylor series for e^x .