

Example 130. Consider the following system of (second-order) initial value problems:

$$\begin{aligned} y_1'' &= 2y_1' - 3y_2' + 7y_2 & y_1(0) &= 2, \quad y_1'(0) = 3, \quad y_2(0) = -1, \quad y_2'(0) = 1 \\ y_2'' &= 4y_1' + y_2' - 5y_1 \end{aligned}$$

Write it as a first-order initial value problem in the form $\mathbf{y}' = A\mathbf{y}$, $\mathbf{y}(0) = \mathbf{y}_0$.

Solution. Introduce $y_3 = y_1'$ and $y_4 = y_2'$. Then, the given system translates into

$$\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 7 & 2 & -3 \\ -5 & 0 & 4 & 1 \end{bmatrix} \mathbf{y}, \quad \mathbf{y}(0) = \begin{bmatrix} 2 \\ -1 \\ 3 \\ 1 \end{bmatrix}.$$

Review. Jordan normal form

Example 131.

- (a) What are the possible Jordan normal forms of a 3×3 matrix with eigenvalues $3, 3, 3$?
- (b) What are the possible Jordan normal forms of a 4×4 matrix with eigenvalues $3, 3, 3, 3$?
- (c) What if the matrix is 5×5 and has eigenvalues $4, 4, 3, 3, 3$?

Solution.

(a) $\begin{bmatrix} 3 & & \\ & 3 & \\ & & 3 \end{bmatrix}, \begin{bmatrix} 3 & 1 & \\ & 3 & 1 & \\ & & 3 & \end{bmatrix}, \begin{bmatrix} 3 & 1 & \\ & 3 & 1 & \\ & & 3 & \end{bmatrix}$

The dimension of the 3-eigenspace equals the number of Jordan blocks: 3, 2, 1, respectively.

Comment. Note that, say, $\begin{bmatrix} 3 & 1 & \\ & 3 & \\ & & 3 \end{bmatrix}$ is equivalent to $\begin{bmatrix} 3 & & \\ & 3 & 1 & \\ & & 3 & \end{bmatrix}$ because the ordering of the diagonal blocks does not matter (as you know from diagonalization).

(b) Now, there are 5 possibilities:

$$\begin{bmatrix} 3 & & & \\ & 3 & & \\ & & 3 & \\ & & & 3 \end{bmatrix}, \begin{bmatrix} 3 & & & \\ & 3 & & \\ & & 3 & 1 & \\ & & & 3 & \end{bmatrix}, \begin{bmatrix} 3 & 1 & & \\ & 3 & & \\ & & 3 & 1 & \\ & & & 3 & \end{bmatrix}, \begin{bmatrix} 3 & & & \\ & 3 & 1 & \\ & & 3 & 1 & \\ & & & 3 & \end{bmatrix}, \begin{bmatrix} 3 & 1 & & \\ & 3 & 1 & \\ & & 3 & 1 & \\ & & & 3 & \end{bmatrix}$$

The dimension of the 3-eigenspace equals the number of Jordan blocks: 4, 3, 2, 2, 1, respectively.

(c) $\begin{bmatrix} 3 & & & & \\ & 3 & & & \\ & & 3 & & \\ & & & 4 & \\ & & & & 4 \end{bmatrix}, \begin{bmatrix} 3 & & & & \\ & 3 & & & \\ & & 3 & & \\ & & & 4 & 1 & \\ & & & & 4 & \end{bmatrix}, \begin{bmatrix} 3 & & & & \\ & 3 & 1 & & \\ & & 3 & & \\ & & & 4 & \\ & & & & 4 \end{bmatrix}, \begin{bmatrix} 3 & & & & \\ & 3 & 1 & & \\ & & 3 & & \\ & & & 4 & 1 & \\ & & & & 4 & \end{bmatrix}, \begin{bmatrix} 3 & 1 & & & \\ & 3 & 1 & & \\ & & 3 & & \\ & & & 4 & \\ & & & & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 & & & \\ & 3 & 1 & & \\ & & 3 & & \\ & & & 4 & 1 & \\ & & & & 4 & \end{bmatrix}$

Note that this is just all possible (namely, 3) Jordan normal forms of a 3×3 matrix with eigenvalues $3, 3, 3$ combined with all possible (namely, 2) Jordan normal forms of a 2×2 matrix with eigenvalues $4, 4$. In total, that makes $3 \cdot 2 = 6$ possibilities.

Comment. Let $p(n)$ be the number of inequivalent Jordan normal forms of an $n \times n$ matrix with a single eigenvalue, n times repeated. We have seen that $p(2) = 2$, $p(3) = 3$, $p(4) = 5$. Note that $p(n)$ is equal to the number of ways of writing n as an ordered sum of positive integers: for instance, $p(4) = 5$ because $4 = 3 + 1 = 2 + 2 = 2 + 1 + 1 = 1 + 1 + 1 + 1$.

$p(n)$ is referred to as the **partition function** and, surprisingly, is a remarkably interesting mathematical object. [https://en.wikipedia.org/wiki/Partition_function_\(number_theory\)](https://en.wikipedia.org/wiki/Partition_function_(number_theory))

Example 132. (summary of small cases)

(a) There are 2 possible Jordan normal forms of a 2×2 matrix with eigenvalues λ, λ .

Namely. $\begin{bmatrix} \lambda & \\ & \lambda \end{bmatrix}, \begin{bmatrix} \lambda & 1 \\ & \lambda \end{bmatrix}$

(b) There are 3 possible Jordan normal forms of a 3×3 matrix with eigenvalues $\lambda, \lambda, \lambda$.

Namely. $\begin{bmatrix} \lambda & & \\ & \lambda & \\ & & \lambda \end{bmatrix}, \begin{bmatrix} \lambda & & \\ & \lambda & 1 \\ & & \lambda \end{bmatrix}, \begin{bmatrix} \lambda & 1 & \\ & \lambda & 1 \\ & & \lambda \end{bmatrix}$

(c) There are 5 possible Jordan normal forms of a 4×4 matrix with eigenvalues $\lambda, \lambda, \lambda, \lambda$.

Namely. $\begin{bmatrix} \lambda & & & \\ & \lambda & & \\ & & \lambda & \\ & & & \lambda \end{bmatrix}, \begin{bmatrix} \lambda & & & \\ & \lambda & & \\ & & \lambda & 1 \\ & & & \lambda \end{bmatrix}, \begin{bmatrix} \lambda & 1 & & \\ & \lambda & & \\ & & \lambda & 1 \\ & & & \lambda \end{bmatrix}, \begin{bmatrix} \lambda & & 1 & \\ & \lambda & 1 & \\ & & \lambda & 1 \\ & & & \lambda \end{bmatrix}, \begin{bmatrix} \lambda & 1 & & \\ & \lambda & 1 & \\ & & \lambda & 1 \\ & & & \lambda \end{bmatrix}$

Example 133. What are the possible Jordan normal forms of a 6×6 matrix with eigenvalues 3, 3, 7, 7, 7, 7?

Solution. There are $2 \cdot 5 = 10$ possible Jordan normal forms for such a matrix:

$$\begin{bmatrix} 3 & & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & 1 & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & 1 & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & 1 & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & 1 & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}, \begin{bmatrix} 3 & 1 & & & & \\ & 3 & & & & \\ & & 7 & & & \\ & & & 7 & & \\ & & & & 7 & 1 \\ & & & & & 7 \end{bmatrix}$$

Example 134. How many different Jordan normal forms are there in the following cases?

- (a) A 8×8 matrix with eigenvalues 1, 1, 2, 2, 2, 4, 4, 4?
- (b) A 11×11 matrix with eigenvalues 1, 1, 1, 2, 2, 2, 2, 4, 4, 4, 4?

Solution.

- (a) $2 \cdot 3 \cdot 3 = 18$ possible Jordan normal forms
- (b) $3 \cdot 5 \cdot 5 = 75$ possible Jordan normal forms