

Review: Matrix calculus

Example 1. Matrix multiplication is not commutative!

$$\bullet \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 3 & 10 \end{bmatrix}$$

Multiplication (on the right) with that “almost identity matrix” is performing the column operation $C_2 + 2C_1 \Rightarrow C_2$ (i.e. 2 times the first column is added to the second column).

$$\bullet \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 3 & 4 \end{bmatrix}$$

Multiplication (on the left) with the same matrix is performing the row operation $R_1 + 2R_2 \Rightarrow R_1$.

First comment. This indicates a second interpretation of matrix multiplication: instead of taking linear combinations of columns of the first matrix, we can also take linear combinations of rows of the second matrix.

Second comment. The row operations we are doing during Gaussian elimination can be realized by multiplying (on the left) with “almost identity matrices”.

Example 2. $\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 14 \end{bmatrix}$ whereas $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{bmatrix}$.

If you know about the dot product, do you see a connection with the first case?

Example 3. Suppose A is $m \times n$ and B is $p \times q$. When does AB make sense? In that case, what are the dimensions of AB ?

AB makes sense if $n = p$. In that case, AB is a $m \times q$ matrix.

Example 4. $\begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

On the RHS we have the **identity matrix**, usually denoted I or I_2 (since it's the 2×2 identity matrix here).

Hence, the two matrices on the left are inverses of each other: $\begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}$, $\begin{bmatrix} 1 & -1 \\ -2 & 3 \end{bmatrix}^{-1} = \begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$.

Example 5. The following formula immediately gives us the inverse of a 2×2 matrix (if it exists). It is worth remembering!

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \quad \text{provided that } ad-bc \neq 0$$

Let's check that! $\frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \frac{1}{ad-bc} \begin{bmatrix} ad-bc & 0 \\ 0 & -cb+ad \end{bmatrix} = I_2$

In particular, a 2×2 matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is invertible $\iff ad-bc \neq 0$.

Recall that this is the **determinant**: $\det\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) = ad-bc$.

In particular:

$$\det(A) = 0 \iff A \text{ is not invertible}$$

Similarly, for $n \times n$ matrices A :

A is invertible	(i.e. there is a matrix A^{-1} such that $AA^{-1} = I$)
$\iff \det(A) \neq 0$	
$\iff Ax = b$ has a unique solution	(namely, $x = A^{-1}b$)

Comment. Why is it not common to write $\frac{1}{A}$ instead of A^{-1} ?

The notation $\frac{1}{A}$ easily leads to ambiguities: for instance, should $\frac{B}{A}$ mean BA^{-1} or should it mean $A^{-1}B$?

[Of course, one could try to avoid this by notations like B/A which would more clearly mean BA^{-1} . It's just not common and doesn't have any real advantages.]

Course comment:

Homework is posted after every class to our course website.

Today's homework needs to be submitted online before 1/20.

Example 6.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -7 & 2 & 3 \\ -16 & 5 & 6 \\ -25 & 8 & 9 \end{bmatrix}$$

Multiplication (on the right) with that “almost identity matrix” is performing the column operation $C_1 - 4C_2 \Rightarrow C_1$ (i.e. -4 times the second column is added to the first column).

$$\begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 7 & 8 & 9 \end{bmatrix}$$

Multiplication (on the left) with the same matrix is performing the row operation $R_2 - 4R_1 \Rightarrow R_2$.

Comment (again). The row operations we are doing during Gaussian elimination can all be realized by multiplying (on the left) with “almost identity matrices”.

These matrices are called **elementary matrices** (they are obtained by performing a single elementary row operation on an identity matrix).

Elementary matrices are **invertible** because elementary row operations are reversible:

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & & \\ & \frac{1}{2} & \\ & & 1 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example 7. Let us do Gaussian elimination on $A = \begin{bmatrix} 2 & 1 \\ 4 & -6 \end{bmatrix}$ until we have an echelon form:

$$A = \begin{bmatrix} 2 & 1 \\ 4 & -6 \end{bmatrix} \xrightarrow{R_2 - 2R_1 \Rightarrow R_2} \begin{bmatrix} 2 & 1 \\ 0 & -8 \end{bmatrix}$$

As last class, the row operation can be encoded by multiplication with an “almost identity matrix” E :

$$\underbrace{\begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}}_E \underbrace{\begin{bmatrix} 2 & 1 \\ 4 & -6 \end{bmatrix}}_A = \underbrace{\begin{bmatrix} 2 & 1 \\ 0 & -8 \end{bmatrix}}_U$$

Since $\begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$ (no calculation needed!), this means that

$$A = E^{-1}U = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 0 & -8 \end{bmatrix}.$$

We factored A as the product of a lower and an upper triangular matrix!

$A = LU$ is known as the **LU decomposition** of A .
 L is lower triangular, U is upper triangular.

If A is $m \times n$, then L is an invertible lower triangular $m \times m$ matrix, and U is a usual **echelon form** of A . Every matrix A has a LU decomposition (after possibly swapping some rows of A first).

- The matrix U is just the echelon form of A produced during Gaussian elimination.
- The matrix L can be constructed, entry-by-entry, by simply recording the row operations used during Gaussian elimination. (No extra work needed!)

Review. The **RREF** (row-reduced echelon form) of A is obtained from an echelon form by

- scaling the pivots to 1, and then
- eliminating the entries above the pivots.

A typical RREF has the shape

[* represents an entry that could be anything]

$$\begin{bmatrix} 1 & * & 0 & * & * & 0 & * \\ & & 1 & * & * & 0 & * \\ & & & & & 1 & * \end{bmatrix}$$

Example 8. The RREF of $A = \begin{bmatrix} 2 & 1 \\ 4 & -6 \end{bmatrix}$ from the previous example is the 2×2 identity matrix.

Comment. That's not surprising: A square matrix is invertible if and only if its RREF is the identity matrix. If that isn't obvious to you, think about how you invert a matrix using Gaussian elimination (reviewed next).

Review. Recall the Gauss–Jordan method of computing A^{-1} . Starting with the augmented matrix $[A \mid I]$, we do Gaussian elimination until we obtain the RREF, which will be of the form $[I \mid A^{-1}]$ so that we can read off A^{-1} .

Why does that work? By our discussion, the steps of Gaussian elimination can be expressed by multiplication (on the left) with a matrix B . Only looking at the first part of the augmented matrix, and since the RREF of an invertible matrix is I , we have $BA = I$, which means that we must have $B = A^{-1}$. The other part of the augmented matrix (which is I initially) gets multiplied with $B = A^{-1}$ as well, so that, in the end, it is $BI = A^{-1}$. That's why we can read off A^{-1} !

For instance. To invert $\begin{bmatrix} 2 & 1 \\ 4 & -6 \end{bmatrix}$ using the Gauss–Jordan method, we would proceed as follows

$$\left[\begin{array}{cc|cc} 2 & 1 & 1 & 0 \\ 4 & -6 & 0 & 1 \end{array} \right] \xrightarrow{R_2 - 2R_1 \Rightarrow R_2} \left[\begin{array}{cc|cc} 2 & 1 & 1 & 0 \\ 0 & -8 & -2 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} \frac{1}{2}R_1 \Rightarrow R_1 \\ -\frac{1}{8}R_2 \Rightarrow R_2 \end{array}} \left[\begin{array}{cc|cc} 1 & \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 1 & \frac{1}{4} & -\frac{1}{8} \end{array} \right] \xrightarrow{R_1 - \frac{1}{2}R_2 \Rightarrow R_1} \left[\begin{array}{cc|cc} 1 & 0 & \frac{3}{8} & \frac{1}{16} \\ 0 & 1 & \frac{1}{4} & -\frac{1}{8} \end{array} \right]$$

and conclude that $\begin{bmatrix} 2 & 1 \\ 4 & -6 \end{bmatrix}^{-1} = \begin{bmatrix} \frac{3}{8} & \frac{1}{16} \\ \frac{1}{4} & -\frac{1}{8} \end{bmatrix}$.

Of course, for 2×2 matrices it is much simpler to use the formula $\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$.

Example 9. (extra) Determine the LU decomposition of $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$.

Solution. $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \xrightarrow{R_2 - 3R_1 \Rightarrow R_2} \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix}$ translates into $\begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix}$.

Since $\begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$ (no calculation needed!), we therefore have $A = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix}$.

Review: Vector spaces, bases, dimension, null spaces

Review.

- Vectors are things that can be **added** and **scaled**.
- Hence, given vectors $\mathbf{v}_1, \dots, \mathbf{v}_n$, the most general we can do is form the **linear combination** $\lambda_1 \mathbf{v}_1 + \dots + \lambda_n \mathbf{v}_n$. The set of all these linear combinations is the **span** of $\mathbf{v}_1, \dots, \mathbf{v}_n$, denoted by $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$.
- Vector **spaces** are spans.

Equivalently. Vector spaces are sets of vectors so that the result of adding and scaling remains within that set.

Homework. Of course, the latter is a very informal statement. Revisit the formal definition, probably consisting of a list of axioms, and observe how that matches with the above (for instance, several of the axioms are concerned with addition and scaling satisfying the “expected” rules).
- Recall that vectors from a vector space V form a **basis** of V if and only if
 - the vectors span V , and
 - the vectors are (linearly) independent.

Equivalently. $\mathbf{v}_1, \dots, \mathbf{v}_n$ from V form a basis of V if and only if every vector in V can be expressed as a unique linear combination of $\mathbf{v}_1, \dots, \mathbf{v}_n$.

Just checking. Make sure that you can define precisely what it means for vectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ to be independent.
- The **dimension** of a vector space V is the number of vectors in a basis for V .
No matter what basis one chooses for V , it always has the same number of vectors.

Example 10. \mathbb{R}^3 is the vector space of all vectors with 3 real entries.

\mathbb{R} itself refers to the set of real numbers. We will later also discuss \mathbb{C} , the set of complex numbers.

The **standard basis** of \mathbb{R}^3 is $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$. The dimension of \mathbb{R}^3 is 3.

Review. The **null space** $\text{null}(A)$ of a matrix A consists of those vectors \mathbf{x} such that $A\mathbf{x} = \mathbf{0}$.

Make sure that you see why $\text{null}(A)$ is a vector space. [For instance, if you pick two vectors in $\text{null}(A)$ why is it that the sum of them is in $\text{null}(A)$ again?]

Example 11. What is $\text{null}(A)$ if the matrix A is invertible?

Solution. If A is invertible, then $A\mathbf{x} = \mathbf{0}$ has the unique solution $\mathbf{x} = A^{-1}\mathbf{0} = \mathbf{0}$.

Hence, $\text{null}(A) = \{\mathbf{0}\}$ which is the trivial vector space (consisting of only the null vector) and has dimension 0.

Example 12. Compute a basis for $\text{null}\left(\begin{bmatrix} -1 & 0 & 2 \\ 2 & -3 & 2 \\ 1 & 0 & -2 \end{bmatrix}\right)$.

Solution. We perform row operations and obtain

$$\text{null}\left(\begin{bmatrix} -1 & 0 & 2 \\ 2 & -3 & 2 \\ 1 & 0 & -2 \end{bmatrix}\right) \stackrel{\substack{R_2+2R_1 \Rightarrow R_2 \\ R_3+R_1 \Rightarrow R_3}}{=} \text{null}\left(\begin{bmatrix} -1 & 0 & 2 \\ 0 & -3 & 6 \\ 0 & 0 & 0 \end{bmatrix}\right) \stackrel{\substack{-R_1 \Rightarrow R_1 \\ -\frac{1}{3}R_2 \Rightarrow R_2}}{=} \text{null}\left(\begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{bmatrix}\right).$$

From the RREF, we can now read off the general solution to $A\mathbf{x} = \mathbf{0}$:

- x_1 and x_2 are pivot variables. [For each we have an equation expressing it in terms of the other variables; for instance, $x_1 - 2x_3 = 0$ tells us that $x_1 = 2x_3$.]
- x_3 is a free variable. [There is no equation forcing a value on x_3 .]

- Hence, without computation, we see that the general solution is $\begin{bmatrix} 2x_3 \\ 2x_3 \\ x_3 \end{bmatrix}$.

In other words, a basis is $\begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$.

Comment. We are starting with the three equations $-x_1 + 2x_3 = 0$, $2x_1 - 3x_2 + 2x_3 = 0$, $x_1 - 2x_3 = 0$. Performing row operations on the matrix is the same as combining these equations (with the objective to form simpler equations by eliminating variables).

Example 13. Compute a basis for $\text{null}\left(\begin{bmatrix} 2 & 0 & 2 \\ 2 & 0 & 2 \\ 1 & 0 & 1 \end{bmatrix}\right)$.

Solution.

$$\text{null}\left(\begin{bmatrix} 2 & 0 & 2 \\ 2 & 0 & 2 \\ 1 & 0 & 1 \end{bmatrix}\right) \stackrel{\substack{R_2-R_1 \Rightarrow R_2 \\ R_3-\frac{1}{2}R_1 \Rightarrow R_3}}{=} \text{null}\left(\begin{bmatrix} 2 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}\right) \stackrel{\frac{1}{2}R_1 \Rightarrow R_1}{=} \text{null}\left(\begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}\right)$$

This time, x_2 and x_3 are free variables. The general solution is $\begin{bmatrix} -x_3 \\ x_2 \\ x_3 \end{bmatrix} = x_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$.

Hence, a basis is $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$.

Review: Eigenvalues and eigenvectors

If $A\mathbf{x} = \lambda\mathbf{x}$ (and $\mathbf{x} \neq \mathbf{0}$), then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ (just a number).

Note that for the equation $A\mathbf{x} = \lambda\mathbf{x}$ to make sense, A needs to be a square matrix (i.e. $n \times n$).

Key observation:

$$\begin{aligned} A\mathbf{x} &= \lambda\mathbf{x} \\ \iff A\mathbf{x} - \lambda\mathbf{x} &= \mathbf{0} \\ \iff (A - \lambda I)\mathbf{x} &= \mathbf{0} \end{aligned}$$

This homogeneous system has a nontrivial solution \mathbf{x} if and only if $\det(A - \lambda I) = 0$.

To find eigenvectors and eigenvalues of A :

(a) First, find the eigenvalues λ by solving $\det(A - \lambda I) = 0$.

$\det(A - \lambda I)$ is a polynomial in λ , called the **characteristic polynomial** of A .

(b) Then, for each eigenvalue λ , find corresponding eigenvectors by solving $(A - \lambda I)\mathbf{x} = \mathbf{0}$.

More precisely, we find a basis of eigenvectors for the λ -**eigenspace** $\text{null}(A - \lambda I)$.

Example 14. $A = \begin{bmatrix} 4 & 0 & 2 \\ 2 & 2 & 2 \\ 1 & 0 & 3 \end{bmatrix}$ has one eigenvector that is “easy” to see. Do you see it?

Solution. Note that $A \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} = 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$. Hence, $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ is a 2-eigenvector.

Just for contrast. Note that $A \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix} \neq \lambda \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$. Hence, $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ is not an eigenvector.

Suppose that A is $n \times n$ and has independent eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_n$.

Then A can be **diagonalized** as $A = PDP^{-1}$, where

- the columns of P are the eigenvectors, and
- the diagonal matrix D has the eigenvalues on the diagonal.

Such a diagonalization is possible if and only if A has enough (independent) eigenvectors.

Comment. If you don't quite recall why these choices result in the diagonalization $A = PDP^{-1}$, note that the diagonalization is equivalent to $AP = PD$.

- Put the eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ as columns into a matrix P .

$$\begin{aligned} A\mathbf{x}_i = \lambda_i\mathbf{x}_i \implies A \begin{bmatrix} | & & | \\ \mathbf{x}_1 & \dots & \mathbf{x}_n \\ | & & | \end{bmatrix} &= \begin{bmatrix} | & & | \\ \lambda_1\mathbf{x}_1 & \dots & \lambda_n\mathbf{x}_n \\ | & & | \end{bmatrix} \\ &= \begin{bmatrix} | & & | \\ \mathbf{x}_1 & \dots & \mathbf{x}_n \\ | & & | \end{bmatrix} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \end{aligned}$$

- In summary: $AP = PD$

Example 15. Let $A = \begin{bmatrix} 4 & 0 & 2 \\ 2 & 2 & 2 \\ 1 & 0 & 3 \end{bmatrix}$.

- (a) Find the eigenvalues and bases for the eigenspaces of A .
- (b) Diagonalize A . That is, determine matrices P and D such that $A = PDP^{-1}$.

Solution.

- (a) By expanding by the second column, we find that the characteristic polynomial $\det(A - \lambda I)$ is

$$\begin{vmatrix} 4-\lambda & 0 & 2 \\ 2 & 2-\lambda & 2 \\ 1 & 0 & 3-\lambda \end{vmatrix} = (2-\lambda) \begin{vmatrix} 4-\lambda & 2 \\ 1 & 3-\lambda \end{vmatrix} = (2-\lambda)[(4-\lambda)(3-\lambda) - 2] = (2-\lambda)^2(5-\lambda).$$

Hence, the eigenvalues are $\lambda = 2$ (with multiplicity 2) and $\lambda = 5$.

Comment. At this point, we know that we will find one eigenvector for $\lambda = 5$ (more precisely, the 5-eigenspace definitely has dimension 1). On the other hand, the 2-eigenspace might have dimension 2 or 1. In order for A to be diagonalizable, the 2-eigenspace must have dimension 2. (Why?!)

- The 5-eigenspace is $\text{null}\left(\begin{bmatrix} -1 & 0 & 2 \\ 2 & -3 & 2 \\ 1 & 0 & -2 \end{bmatrix}\right)$. Proceeding as in Example 12, we obtain

$$\text{null}\left(\begin{bmatrix} -1 & 0 & 2 \\ 2 & -3 & 2 \\ 1 & 0 & -2 \end{bmatrix}\right) \stackrel{\text{RREF}}{=} \text{null}\left(\begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{bmatrix}\right) = \text{span}\left\{\begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}\right\}.$$

In other words, the 5-eigenspace has basis $\begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$.

- The 2-eigenspace is $\text{null}\left(\begin{bmatrix} 2 & 0 & 2 \\ 2 & 0 & 2 \\ 1 & 0 & 1 \end{bmatrix}\right)$. Proceeding as in Example 13, we obtain

$$\text{null}\left(\begin{bmatrix} 2 & 0 & 2 \\ 2 & 0 & 2 \\ 1 & 0 & 1 \end{bmatrix}\right) \stackrel{\text{RREF}}{=} \text{null}\left(\begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}\right) = \text{span}\left\{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}\right\}$$

In other words, the 2-eigenspace has basis $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$.

Comment. So, indeed, the 2-eigenspace has dimension 2. In particular, A is diagonalizable.

- (b) A possible choice is $P = \begin{bmatrix} 2 & 0 & -1 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$, $D = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

Comment. However, many other choices are possible and correct. For instance, the order of the eigenvalues in D doesn't matter (as long as the same order is used for P). Also, for P , the columns can be chosen to be any other set of eigenvectors.

Example 16. (extra practice) Diagonalize, if possible, the matrices

$$A = \begin{bmatrix} 3 & 4 & 1 \\ 0 & 2 & 0 \\ 1 & 4 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Solution. For instance, $A = PDP^{-1}$ with $P = \begin{bmatrix} 1 & -4 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$ and $D = \begin{bmatrix} 4 & & \\ & 2 & \\ & & 2 \end{bmatrix}$. B is not diagonalizable.

For instance, $C = PDP^{-1}$ with $P = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $D = \begin{bmatrix} 1 & & \\ & 0 & \\ & & 0 \end{bmatrix}$.

Example 17. (review) If $A = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$, then its **transpose** is $A^T = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$.

Recall that $(AB)^T = B^T A^T$. This reflects the fact that, in the column-centric versus the row-centric interpretation of matrix multiplication, the order of the matrices is reversed.

Comment. When working with complex numbers, the fundamental role is not played by the transpose but by the **conjugate transpose** instead (we'll see that in our discussion of orthogonality): $A^* = \overline{A^T}$.

For instance, if $A = \begin{bmatrix} 1-3i & 5i \\ 2+i & 3 \end{bmatrix}$, then $A^* = \begin{bmatrix} 1+3i & 2-i \\ -5i & 3 \end{bmatrix}$.

Orthogonality

The inner product and distances

Definition 18. The **inner product** (or **dot product**) of \mathbf{v} , \mathbf{w} in \mathbb{R}^n :

$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \dots + v_n w_n.$$

Because we can think of this as a special case of the matrix product, it satisfies the basic rules like associativity and distributivity.

In addition: $\mathbf{v} \cdot \mathbf{w} = \mathbf{w} \cdot \mathbf{v}$.


Example 19. $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} = 2 - 2 + 12 = 12$

Definition 20.

- The **norm** (or **length**) of a vector \mathbf{v} in \mathbb{R}^n is

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + \dots + v_n^2}.$$

- The **distance** between points \mathbf{v} and \mathbf{w} in \mathbb{R}^n is $\text{dist}(\mathbf{v}, \mathbf{w}) = \|\mathbf{v} - \mathbf{w}\|$.



Example 21. For instance, in \mathbb{R}^2 , $\text{dist}\left(\begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix}\right) = \left\| \begin{bmatrix} x_1 - x_2 \\ y_1 - y_2 \end{bmatrix} \right\| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$.

Example 22. Write $\|\mathbf{v} - \mathbf{w}\|^2$ as a dot product, and multiply it out.

Solution. $\|\mathbf{v} - \mathbf{w}\|^2 = (\mathbf{v} - \mathbf{w}) \cdot (\mathbf{v} - \mathbf{w}) = \mathbf{v} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{w} - \mathbf{w} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = \|\mathbf{v}\|^2 - 2\mathbf{v} \cdot \mathbf{w} + \|\mathbf{w}\|^2$

Comment. This is a vector version of $(x - y)^2 = x^2 - 2xy + y^2$.

The reason we were careful and first wrote $-\mathbf{v} \cdot \mathbf{w} - \mathbf{w} \cdot \mathbf{v}$ before simplifying it to $-2\mathbf{v} \cdot \mathbf{w}$ is that we should not take rules such as $\mathbf{v} \cdot \mathbf{w} = \mathbf{w} \cdot \mathbf{v}$ for granted. For instance, for the cross product $\mathbf{v} \times \mathbf{w}$, that you may have seen in Calculus, we have $\mathbf{v} \times \mathbf{w} \neq \mathbf{w} \times \mathbf{v}$ (instead, $\mathbf{v} \times \mathbf{w} = -\mathbf{w} \times \mathbf{v}$).

Orthogonal vectors

Definition 23. \mathbf{v} and \mathbf{w} in \mathbb{R}^n are **orthogonal** if

$$\mathbf{v} \cdot \mathbf{w} = 0.$$

Why? How is this related to our understanding of right angles?


Pythagoras!

\mathbf{v} and \mathbf{w} are orthogonal

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \underbrace{\|\mathbf{v} - \mathbf{w}\|^2}_{= \|\mathbf{v}\|^2 - 2\mathbf{v} \cdot \mathbf{w} + \|\mathbf{w}\|^2 \text{ (by previous example)}}$$

$$\iff -2\mathbf{v} \cdot \mathbf{w} = 0$$

$$\iff \mathbf{v} \cdot \mathbf{w} = 0$$



Example 24. Determine a basis for the **orthogonal complement** of (the span of) $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$.

What are we looking for? The orthogonal complement of \mathbf{v} consists of all vectors that are orthogonal to \mathbf{v} . More generally, the orthogonal complement of a space V consists of all vectors that are orthogonal to every vector in V .

Solution. (staring/intuition) We are working in 3-dimensional space and already have 1 vector. The vectors orthogonal to it lie in a $3 - 1 = 2$ -dimensional space (a plane).

Two of the vectors orthogonal to $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$ are $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$.

Knowing that the orthogonal complement has dimension 2, we conclude that $\left\{ \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \right\}$ is a basis.

In other words, the orthogonal complement of $\text{span}\left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \right\}$ is $\text{span}\left\{ \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \right\}$.

[Note how the dimensions add up to the dimension of the entire space: $1 + 2 = 3$.]

Solution. (professional) $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$ (dot product!) is the same as $[1 \ 2 \ 1] \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$ (matrix product!).

Hence, the orthogonal complement of $\text{span}\left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \right\}$ is the same as $\text{null}([1 \ 2 \ 1])$.

Computing a basis for $\text{null}([1 \ 2 \ 1])$ is easy since $[1 \ 2 \ 1]$ is already in RREF.

Note that the general solution to $[1 \ 2 \ 1]\mathbf{x} = 0$ is $\begin{bmatrix} -2s - t \\ s \\ t \end{bmatrix} = s \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$.

A basis for $\text{null}([1 \ 2 \ 1])$ therefore is $\left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$. (Check that these are indeed orthogonal to $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$!)

Example 25. Determine a basis for the orthogonal complement of $\text{span}\left\{\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}\right\}$.

Solution. We are looking for vectors \mathbf{x} such that $\begin{bmatrix} 1 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$ and $\begin{bmatrix} 3 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$.

The two equations can be combined into a single one: $\begin{bmatrix} 1 & 2 & 1 \\ 3 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \mathbf{0}$.

In other words, the orthogonal complement of $\text{span}\left\{\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}\right\}$ is the same as $\text{null}\left(\begin{bmatrix} 1 & 2 & 1 \\ 3 & 1 & 2 \end{bmatrix}\right)$.

It remains to compute a basis for that null space:

$$\begin{bmatrix} 1 & 2 & 1 \\ 3 & 1 & 2 \end{bmatrix} \xrightarrow{R_2 - 3R_1} \begin{bmatrix} 1 & 2 & 1 \\ 0 & -5 & -1 \end{bmatrix} \xrightarrow{\text{back-substitution}} \begin{bmatrix} -3/5s \\ -1/5s \\ s \end{bmatrix}$$

Hence, a basis for the orthogonal complement of $\text{span}\left\{\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}\right\}$ is $\begin{bmatrix} -3/5 \\ -1/5 \\ 1 \end{bmatrix}$.

Check. $\begin{bmatrix} -3/5 \\ -1/5 \\ 1 \end{bmatrix}$ is indeed orthogonal to both $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$.

Just to make sure. Why was it clear that the orthogonal complement is 1-dimensional?

The fundamental theorem

Example 26. The four **fundamental subspaces** associated with a matrix A are

$$\text{col}(A), \quad \text{row}(A), \quad \text{null}(A), \quad \text{null}(A^T).$$

Note that $\text{row}(A) = \text{col}(A^T)$. (In particular, we usually write vectors in $\text{row}(A)$ as column vectors.)

Definition 27. $\text{null}(A^T)$ is the **left null space** of A .

Why that name? Recall that, by definition \mathbf{x} is in $\text{null}(A) \iff A\mathbf{x} = \mathbf{0}$.

Likewise, \mathbf{x} is in $\text{null}(A^T) \iff A^T\mathbf{x} = \mathbf{0} \iff \mathbf{x}^T A = \mathbf{0}$.

[Recall that $(AB)^T = B^T A^T$. In particular, $(A^T \mathbf{x})^T = \mathbf{x}^T A$, which is what we used in the last equivalence.]

Theorem 28. (Fundamental Theorem of Linear Algebra, Part I)

Let A be an $m \times n$ matrix of **rank** r .

- $\dim \text{col}(A) = r$ (subspace of \mathbb{R}^m)
- $\dim \text{row}(A) = r$ (subspace of \mathbb{R}^n) $\text{row}(A) = \text{col}(A^T)$
- $\dim \text{null}(A) = n - r$ (subspace of \mathbb{R}^n)
- $\dim \text{null}(A^T) = m - r$ (subspace of \mathbb{R}^m)

Example 29. Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$. Determine bases for all four fundamental subspaces.

Solution. Make sure that, for such a simple matrix, you can see all of these that at a glance!

$$\text{col}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \right\}, \quad \text{row}(A) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}, \quad \text{null}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\}, \quad \text{null}(A^T) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Important observation. The basis vectors for $\text{row}(A)$ and $\text{null}(A)$ are orthogonal! $\begin{bmatrix} -2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \end{bmatrix} = 0$

The same is true for the basis vectors for $\text{col}(A)$ and $\text{null}(A^T)$: $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} = 0$ and $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} -3 \\ 0 \\ 1 \end{bmatrix} = 0$

Vectors in $\text{null}(A)$ are orthogonal to vectors in $\text{row}(A)$.
 In short, $\text{null}(A)$ is orthogonal to $\text{row}(A)$.

Why? Suppose that \mathbf{x} is in $\text{null}(A)$. That is, $A\mathbf{x} = \mathbf{0}$. But think about what $A\mathbf{x} = \mathbf{0}$ means (row-product rule). It means that the inner product of every row with \mathbf{x} is zero. Which implies that \mathbf{x} is orthogonal to the row space.

Definition 30. As done in the observation above, we say that two subspaces V and W of \mathbb{R}^n are **orthogonal** if and only if every vector in V is orthogonal to every vector in W .

The **orthogonal complement** of W is the space W^\perp of all vectors that are orthogonal to W .

Exercise. Show that the orthogonal complement is indeed a vector space.

Theorem 31. (Fundamental Theorem of Linear Algebra, Part II)

- $\text{null}(A)$ is orthogonal to $\text{row}(A)$. (both subspaces of \mathbb{R}^n)

Note that $\dim \text{null}(A) + \dim \text{row}(A) = n$. Hence, the two spaces are orthogonal complements.

- $\text{null}(A^T)$ is orthogonal to $\text{col}(A)$.

Again, the two spaces are orthogonal complements. (This is just the first part with A replaced by A^T .)

Example 32. Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$. Check that $\text{null}(A)$ and $\text{row}(A)$ are orthogonal complements.

Solution. $\begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix} \xrightarrow[\begin{smallmatrix} R_2 - 2R_1 \Rightarrow R_2 \\ R_3 - 3R_1 \Rightarrow R_3 \end{smallmatrix}]{\rightsquigarrow} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 0 & -2 \\ 0 & 0 & -3 \end{bmatrix} \xrightarrow[\rightsquigarrow]{R_3 - \frac{3}{2}R_2 \Rightarrow R_3} \begin{bmatrix} 1 & 2 & 1 \\ 0 & 0 & -2 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow[\rightsquigarrow]{\text{RREF}} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$

Hence, $\text{null}(A) = \text{span}\left\{\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}\right\}$, $\text{row}(A) = \text{span}\left\{\begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right\}$.

$\text{null}(A)$ and $\text{row}(A)$ are orthogonal because

$$\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = 0, \quad \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0.$$

Note. Because $\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$ is orthogonal to both basis vectors, it is orthogonal to every vector in $\text{row}(A)$.

Vectors in $\text{row}(A)$ are of the form $\mathbf{v} = a \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} + b \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$. Then, $\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \mathbf{v} = a \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} + b \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0$.

Conclusion. Hence, $\text{null}(A)$ and $\text{row}(A)$ are indeed orthogonal spaces.

In fact, $\text{null}(A)$ and $\text{row}(A)$ are orthogonal complements.

That is because $\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ are orthogonal, hence independent, and thus a basis of all of \mathbb{R}^3 .

Example 33. (extra) Determine bases for all four fundamental subspaces of

$$A = \begin{bmatrix} 1 & 2 & 1 & 3 \\ 2 & 4 & 0 & 1 \\ 3 & 6 & 0 & 1 \end{bmatrix}.$$

Verify all parts of the Fundamental Theorem, especially that $\text{null}(A)$ and $\text{row}(A)$ (as well as $\text{null}(A^T)$ and $\text{col}(A)$) are orthogonal complements.

Partial solution. One can almost see that $\text{rank}(A) = 3$. Hence, the dimensions of the fundamental subspaces are ...

Any serious linear algebra problems are done by a machine. Let us see how to use the open-source computer algebra system **Sage** to do basic computations for us.

Sage is freely available at sagemath.org. Instead of installing it locally (it's huge!) we can conveniently use it in the cloud at cocalc.com from any browser. For short computations, like the one below, you can also just use the input field on our course website.

Sage is built as a **Python** library, so any Python code is valid. Here, we will just use it as a fancy calculator.

Let's revisit Example 32 and let Sage do the work for us:

```
Sage] A = matrix([[1,2,1],[2,4,0],[3,6,0]])
```

```
Sage] A.rref()
```

$$\begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Similarly, if we wanted to compute a basis for $\text{null}(A^T)$, we can simply do:

```
Sage] A.transpose().rref()
```

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & \frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix}$$

Here's some other standard things we might be interested in (compare with Example 15):

```
Sage] A = matrix([[4,0,2],[2,2,2],[1,0,3]])
```

```
Sage] A.eigenvalues()
```

```
[5, 2, 2]
```

```
Sage] A.eigenvectors_right()
```

$$\left[\left(5, \left[\left(1, 1, \frac{1}{2} \right) \right], 1 \right), \left(2, \left[(1, 0, -1), (0, 1, 0) \right], 2 \right) \right]$$

```
Sage] A.eigenmatrix_right()
```

$$\left(\begin{bmatrix} 5 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ \frac{1}{2} & -1 & 0 \end{bmatrix} \right)$$

```
Sage] A.rank()
```

```
3
```

```
Sage] A.determinant()
```

```
20
```

```
Sage] A.inverse()
```

$$\begin{bmatrix} \frac{3}{10} & 0 & -\frac{1}{5} \\ -\frac{1}{5} & \frac{1}{2} & -\frac{1}{5} \\ -\frac{1}{10} & 0 & \frac{1}{5} \end{bmatrix}$$

Example 34. (warmup) $\begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \\ 5 \end{bmatrix}$

Note that this means that the system of equations $\begin{matrix} x_1 + 2x_2 = 1 \\ 3x_1 + x_2 = 1 \\ 5x_2 = 1 \end{matrix}$ can also be written as $\begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$.

[This was the motivation for introducing matrix-vector multiplication.]

In the same way, any system can be written as $Ax = b$, where A is a matrix and b a vector. In particular, this makes it obvious that:

$$Ax = b \text{ is consistent} \iff b \text{ is in } \text{col}(A)$$

Recall that, by the FTLA, $\text{col}(A)$ and $\text{null}(A^T)$ are orthogonal complements.

Theorem 35. $Ax = b$ is consistent $\iff b$ is orthogonal to $\text{null}(A^T)$

Proof. $Ax = b$ is consistent $\iff b$ is in $\text{col}(A) \xleftrightarrow{\text{FTLA}} b$ is orthogonal to $\text{null}(A^T)$

Note. b is orthogonal to $\text{null}(A^T)$ means that $y^T b = 0$ whenever $y^T A = 0$. Why?!

Example 36. Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix}$. For which b does $Ax = b$ have a solution?

Solution. (old)

$$\left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 3 & 1 & b_2 \\ 0 & 5 & b_3 \end{array} \right] \xrightarrow{R_2 - 3R_1 \Rightarrow R_2} \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & -5 & -3b_1 + b_2 \\ 0 & 5 & b_3 \end{array} \right] \xrightarrow{R_3 + R_2 \Rightarrow R_3} \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & -5 & -3b_1 + b_2 \\ 0 & 0 & -3b_1 + b_2 + b_3 \end{array} \right]$$

So, $Ax = b$ is consistent if and only if $-3b_1 + b_2 + b_3 = 0$.

Solution. (new) We determine a basis for $\text{null}(A^T)$:

$$\left[\begin{array}{ccc} 1 & 3 & 0 \\ 2 & 1 & 5 \end{array} \right] \xrightarrow{R_2 - 2R_1 \Rightarrow R_2} \left[\begin{array}{ccc} 1 & 3 & 0 \\ 0 & -5 & 5 \end{array} \right] \xrightarrow{-\frac{1}{5}R_2 \Rightarrow R_2} \left[\begin{array}{ccc} 1 & 3 & 0 \\ 0 & 1 & -1 \end{array} \right] \xrightarrow{R_1 - 3R_2 \Rightarrow R_1} \left[\begin{array}{ccc} 1 & 0 & 3 \\ 0 & 1 & -1 \end{array} \right]$$

We read off from the RREF that $\text{null}(A^T)$ has basis $\begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$.

b has to be orthogonal to $\text{null}(A^T)$. That is, $b \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} = 0$. As above!

Comment. Below is how we can use Sage to (try and) solve $Ax = b$ for $b = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ and $b = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$.

```
Sage] A = matrix([[1,2],[3,1],[0,5]])
```

```
Sage] A.solve_right(vector([1,1,2]))
```

$$\left(\frac{1}{5}, \frac{2}{5} \right)$$

```
Sage] A.solve_right(vector([1,1,1]))
```

```
Traceback (most recent call last):
ValueError: matrix equation has no solutions
```


Least squares

Example 37. Not all linear systems have solutions.


In fact, for many applications, data needs to be fitted and there is no hope for a perfect match.

For instance, $Ax = b$ with

$$\begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix} x = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

has no solution:

- $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ is not in $\text{col}(A)$ since $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} \neq 0$ (see previous example).
- Instead of giving up, we want the x which makes Ax and b as close as possible.
- Such x is characterized by the error $Ax - b$ being **orthogonal** to $\text{col}(A)$ (i.e. all possible Ax).



Definition 38. \hat{x} is a **least squares solution** of the system $Ax = b$ if \hat{x} is such that $A\hat{x} - b$ is as small as possible (i.e. minimal norm).

- If $Ax = b$ is consistent, then \hat{x} is just an ordinary solution. (in that case, $A\hat{x} - b = 0$)
- Interesting case: $Ax = b$ is inconsistent. (in particular, if the system is overdetermined)

The normal equations

The following result provides a straightforward recipe (thanks to the FTLA) to find least squares solutions for any system $Ax = b$.

Theorem 39. \hat{x} is a least squares solution of $Ax = b$
 $\iff A^T A \hat{x} = A^T b$ (the **normal equations**)

Proof.

\hat{x} is a least squares solution of $Ax = b$

$\iff A\hat{x} - b$ is as small as possible

$\iff A\hat{x} - b$ is orthogonal to $\text{col}(A)$

$\stackrel{\text{FTLA}}{\iff} A\hat{x} - b$ is in $\text{null}(A^T)$

$\iff A^T(A\hat{x} - b) = 0$

$\iff A^T A \hat{x} = A^T b$

□

Example 40. Find the least squares solution to $Ax = b$, where

$$A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}.$$

Solution. First, $A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$ and $A^T b = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

Hence, the normal equations $A^T A \hat{x} = A^T b$ take the form $\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \hat{x} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

Solving, we immediately find $\hat{x} = \begin{bmatrix} 1/2 \\ 3/2 \end{bmatrix}$.

Check. Since $A\hat{x} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$, the error is $A\hat{x} - b = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}$. Recall that the error must be orthogonal to $\text{col}(A)$!

This error is indeed orthogonal to $\text{col}(A)$ because $\begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = 0$ and $\begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = 0$.

Comment. Why are the normal equations so particularly simple (compare with example below for the typical case) here? Note how each entry of the product $A^T A$ is computed as the dot product of two columns of A (matrix products of a row of A^T times a column of A). That $A^T A$ is a diagonal matrix reflects the fact that the two columns of A are orthogonal to each other.

Example 41. (extra) Find the least squares solution to $Ax = b$, where

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

Solution. First, $A^T A = \begin{bmatrix} 1 & 3 & 0 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} 10 & 5 \\ 5 & 30 \end{bmatrix}$ and $A^T b = \begin{bmatrix} 1 & 3 & 0 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$.

Hence, the normal equations $A^T A \hat{x} = A^T b$ take the form $\begin{bmatrix} 10 & 5 \\ 5 & 30 \end{bmatrix} \hat{x} = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$.

Since $\begin{bmatrix} 10 & 5 \\ 5 & 30 \end{bmatrix}^{-1} = \frac{1}{275} \begin{bmatrix} 30 & -5 \\ -5 & 10 \end{bmatrix} = \frac{1}{55} \begin{bmatrix} 6 & -1 \\ -1 & 2 \end{bmatrix}$, we find $\hat{x} = \frac{1}{55} \begin{bmatrix} 6 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 4 \\ 8 \end{bmatrix} = \frac{1}{55} \begin{bmatrix} 16 \\ 12 \end{bmatrix}$.

Check. Since $A\hat{x} = \frac{1}{55} \begin{bmatrix} 40 \\ 60 \\ 60 \end{bmatrix}$, the error $A\hat{x} - b = \frac{1}{55} \begin{bmatrix} -15 \\ 5 \\ 5 \end{bmatrix} = \frac{1}{11} \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}$ must be orthogonal to $\text{col}(A)$.

The error is indeed orthogonal to $\text{col}(A)$ because $\begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix} \cdot \frac{1}{11} \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} = 0$ and $\begin{bmatrix} 2 \\ 1 \\ 5 \end{bmatrix} \cdot \frac{1}{11} \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix} = 0$.

Example 42. Find the least squares solution to $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}.$$

Solution. First, $A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$ and $A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Hence, the normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ take the form $\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Solving, we immediately find $\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

Check. The error $A\hat{\mathbf{x}} - \mathbf{b} = \begin{bmatrix} 2 \\ 4 \\ -8 \end{bmatrix}$ is indeed orthogonal to $\text{col}(A)$. Because $\begin{bmatrix} 2 \\ 4 \\ -8 \end{bmatrix} \cdot \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix} = 0$ and $\begin{bmatrix} 2 \\ 4 \\ -8 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix} = 0$.

Application: least squares lines

Given data points (x_i, y_i) , we wish to find optimal parameters a, b such that $y_i \approx a + bx_i$ for all i .

Example 43. Determine the line that “best fits” the data points $(2, 1), (5, 2), (7, 3), (8, 3)$.

Comment. Can you see that there is no line fitting the data perfectly? (Check out the last two points!)

Solution. We need to determine the values a, b for the best-fitting line $y = a + bx$.

If there was a line that fit the data perfectly, then:

$$\begin{aligned} a + 2b &= 1 && (2, 1) \\ a + 5b &= 2 && (5, 2) \\ a + 7b &= 3 && (7, 3) \\ a + 8b &= 3 && (8, 3) \end{aligned}$$

In matrix form, this is: $\underbrace{\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}}_{\text{design matrix } X} \begin{bmatrix} a \\ b \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}}_{\text{observation vector } \mathbf{y}}$ (writing the points as (x_i, y_i))

Using our points, these equations become $\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$. [This system is inconsistent (as expected).]

We compute a least squares solution.

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}, \quad X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}.$$

Solving the normal equations $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$.

Hence, the least squares line is $y = \frac{2}{7} + \frac{5}{14}x$.

The plot above shows our points together with this line. It does look like a very good fit!

Important comment. In what sense is this the line of “best fit”? By computing a least squares solution the way we do, we are minimizing the error $\mathbf{y} - X \begin{bmatrix} a \\ b \end{bmatrix}$. The components of that error are $y_i - (a + bx_i)$.

Hence, we see that we are minimizing the **residual sum of squares** $SS_{\text{res}} = \sum_i [y_i - (a + bx_i)]^2$.

Also see the discussion after the next example (where we swap the role of x and y) as well as the example at the beginning of next class (where we discuss making predictions and why minimizing SS_{res} corresponds to minimizing the error of those predictions).

Example 44. (again) Determine the least squares line for the points $(2, 1)$, $(5, 2)$, $(7, 3)$, $(8, 3)$.

Solution. Let's repeat the computation we did last class. This time, we let Sage do the actual work for us:

```
Sage] X = matrix([[1,2],[1,5],[1,7],[1,8]]); y = vector([1,2,3,3])
```

```
Sage] (X.transpose()*X).solve_right(X.transpose()*y)
```

$$\left(\frac{2}{7}, \frac{5}{14}\right)$$

Here are some intermediate steps to help see what's going on (and that it matches our earlier work):

```
Sage] X.transpose()*X
```

$$\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$


```
Sage] X.transpose()*y
```

$$(9, 57)$$


Let's plot the least squares line $y = \frac{2}{7} + \frac{5}{14}x$ in Sage to marvel at the good fit!

```
Sage] points = [[2,1],[5,2],[7,3],[8,3]]
```


```
Sage] scatter_plot(points)
```



```
Sage] scatter_plot(points) + plot(2/7+5/14*x,1,9)
```



Comment. As mentioned earlier, the least squares line minimizes the (sum of squares of the) vertical offsets:



<http://mathworld.wolfram.com/LeastSquaresFitting.html>

Comment. We get a (slightly) different “best fit” line if we change the role of x and y ! Can you explain that?


```
Sage] X = matrix([[1,1],[1,2],[1,3],[1,3]]); y = vector([2,5,7,8])
```

```
Sage] (X.transpose()*X).solve_right(X.transpose()*y)
```

$$\left(-\frac{7}{11}, \frac{30}{11}\right)$$

Note that $x = -\frac{7}{11} + \frac{30}{11}y$ is equivalent to $y = \frac{7}{30} + \frac{11}{30}x$.

```
Sage] scatter_plot([[2,1],[5,2],[7,3],[8,3]]) + plot(2/7+5/14*x,1,9) + plot(7/30+11/30*x,
1,9,color='red')
```



The explanation is that (see pictures at the beginning of this example) we are minimizing vertical offsets in one case and horizontal offsets in the other case.

In linear regression, the relationship between a dependent variable and one or more explanatory variables is modeled. If y is the dependent variable, with x the explanatory variable, then it is natural to minimize the error we make in “predicting y through x ” (vertical offsets). See example at the beginning of next class!

Example 45. A car rental company wants to predict the annual maintenance cost y (in 100USD/year) of a car using the age x (in years) of that car (as an explanatory variable). Based on the observations $(x, y) = (2, 1), (5, 2), (7, 3), (8, 3)$, predict the cost for a 4.5 year old car (using linear regression).

Solution. Once we compute the regression line $y = a + bx$ (we already did that: $y = \frac{2}{7} + \frac{5}{14}x$), our prediction is $\frac{2}{7} + \frac{5}{14} \cdot 4.5 = \frac{53}{28} \approx 1.89$, that is, 189 USD/year.

Application: multiple linear regression

In statistics, **linear regression** is an approach for modeling the relationship between a scalar dependent variable and one or more explanatory variables.

The case of one explanatory variable is called simple linear regression.

For more than one explanatory variable, the process is called multiple linear regression.

http://en.wikipedia.org/wiki/Linear_regression

The experimental data might be of the form (x_i, y_i, z_i) , where now the dependent variable z_i depends on two explanatory variables x_i, y_i (instead of just x_i).

Example 46. Set up a linear system to find values for the parameters a, b, c such that $z = a + bx + cy$ best fits some given points $(x_1, y_1, z_1), (x_2, y_2, z_2), \dots$

Solution. The equations $a + bx_i + cy_i = z_i$ translate into the system:

$$\underbrace{\begin{bmatrix} 1 & x_1 & y_1 \\ 1 & x_2 & y_2 \\ 1 & x_3 & y_3 \\ \vdots & \vdots & \vdots \end{bmatrix}}_{\text{design matrix } A} \underbrace{\begin{bmatrix} a \\ b \\ c \end{bmatrix}}_{\text{observation vector } \mathbf{z}} = \underbrace{\begin{bmatrix} z_1 \\ z_2 \\ z_3 \\ \vdots \end{bmatrix}}_{\text{observation vector } \mathbf{z}}$$

Of course, this is usually inconsistent. To find the best possible a, b, c we compute a least squares solution by solving $A^T A \begin{bmatrix} a \\ b \\ c \end{bmatrix} = A^T \mathbf{z}$.

Application: Fitting data to other curves

We can also fit the experimental data (x_i, y_i) using other curves.

Example 47. Set up a linear system to find values for the parameters a, b, c that result in the quadratic curve $y = a + bx + cx^2$ that best fits some given points $(x_1, y_1), (x_2, y_2), \dots$

Solution. $y_i \approx a + bx_i + cx_i^2$ with parameters a, b, c .

The equations $y_i = a + bx_i + cx_i^2$ in matrix form:

$$\underbrace{\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ \vdots & \vdots & \vdots \end{bmatrix}}_{\text{design matrix } A} \underbrace{\begin{bmatrix} a \\ b \\ c \end{bmatrix}}_{\text{observation vector } \mathbf{y}} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \end{bmatrix}}_{\text{observation vector } \mathbf{y}}$$

Again, we determine values for a, b, c by computing a least squares solution to that system.

That is, we need to solve the system $A^T A \begin{bmatrix} a \\ b \\ c \end{bmatrix} = A^T \mathbf{y}$.

Example 48. (extra) Use Sage to find values for a, b, c that result in the quadratic curve $y = a + bx + cx^2$ that best fits the points $(0, 1), (1, 2), (2, 3), (3, -4), (4, -7), (5, -12)$.

Solution. We first input the points:

```
Sage] points = [[0,1],[1,2],[2,3],[3,-4],[4,-7],[5,-12]]
```

We set up the system described in the previous example, then determine a least-squares solution.

```
Sage] X = matrix([[1,0,0],[1,1,1],[1,2,4],[1,3,9],[1,4,16],[1,5,25]])
```


```
Sage] y = vector([1,2,3,-4,-7,-12])
```

```
Sage] (X.transpose()*X).solve_right(X.transpose()*y)
```

$$\left(\frac{3}{2}, \frac{179}{140}, -\frac{23}{28} \right)$$

Hence, the best fitting quadratic curve is $y = \frac{3}{2} + \frac{179}{140}x - \frac{23}{28}x^2$. Here's a plot:

```
Sage] scatter_plot(points) + plot(3/2+179/140*x-23/28*x^2,0,5,color='red')
```



Advanced comment. If you are comfortable with Python, you can avoid typing out X and \mathbf{y} :
[The plot command above now won't work anymore because we are overwriting x with numbers.]

```
Sage] X = matrix([[1,x,x^2] for x,y in points])
```

```
Sage] y = vector([y for x,y in points])
```

More on orthogonality

Projection matrices

The **(orthogonal) projection** $\hat{\mathbf{b}}$ of a vector \mathbf{b} onto a subspace W is the vector in W closest to \mathbf{b} .

We can compute $\hat{\mathbf{b}}$ as follows:

- Write $W = \text{col}(A)$ for some matrix A .
- Then $\hat{\mathbf{b}} = A\hat{\mathbf{x}}$ where $\hat{\mathbf{x}}$ is a least squares solution to $A\mathbf{x} = \mathbf{b}$. (i.e. $\hat{\mathbf{x}}$ solves $A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$)

Why? Why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{col}(A)$?

Because, for a least squares solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible (and any element in $\text{col}(A)$ is of the form $A\mathbf{x}$ for some \mathbf{x}).

Note. This is a recipe for computing any orthogonal projection! That's because every subspace W can be written as $\text{col}(A)$ for some choice of the matrix A (take, for instance, A so that its columns are a basis for W).

Example 49. What is the orthogonal projection of $\begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$ onto $\text{span}\left\{\begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}\right\}$?

Solution. In other words, what is the orthogonal projection of $\mathbf{b} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$ onto $\text{col}(A)$ with $A = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}$.

In Example 42, we found that the system $A\mathbf{x} = \mathbf{b}$ has the least squares solution $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

The projection $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{col}(A)$ thus is $A\hat{\mathbf{x}} = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 3 \end{bmatrix}$.

Check. The error $\hat{\mathbf{b}} - \mathbf{b} = \begin{bmatrix} 2 \\ 4 \\ -8 \end{bmatrix}$ needs to be orthogonal to $\text{col}(A)$. Indeed: $\begin{bmatrix} 2 \\ 4 \\ -8 \end{bmatrix} \cdot \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix} = 0$ and $\begin{bmatrix} 2 \\ 4 \\ -8 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix} = 0$.

Example 50. (extra)

(a) What is the orthogonal projection of $\begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$ onto $\text{span}\left\{\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}\right\}$?

(b) What is the orthogonal projection of $\begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$ onto $\text{span}\left\{\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}\right\}$?

Solution. (final answer only) The projections are $\left(\frac{11}{6}, \frac{1}{3}, \frac{7}{6}\right)^T$ and $\left(\frac{3}{2}, 0, \frac{3}{2}\right)^T$.

Review. We can compute the orthogonal projection $\hat{\mathbf{b}}$ of \mathbf{b} onto W as follows:

- Write $W = \text{col}(A)$, where the columns of A are a basis of W .
Then, $\hat{\mathbf{b}} = A\hat{\mathbf{x}}$ where $\hat{\mathbf{x}}$ is the least squares solution to $A\mathbf{x} = \mathbf{b}$ (i.e. $A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$).

Assuming $A^T A$ is invertible (which, as discussed in the lemma below, is automatically the case if the columns of A are independent), we have $\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b}$ and hence:

(projection matrix) The projection $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{col}(A)$ is (assuming cols of A are independent)

$$\hat{\mathbf{b}} = \underbrace{A(A^T A)^{-1} A^T}_{P} \mathbf{b}.$$

The matrix $P = A(A^T A)^{-1} A^T$ is the **projection matrix** for projecting onto $\text{col}(A)$.

Lemma 51. If the columns of a matrix A are independent, then $A^T A$ is invertible.

Proof. Assume $A^T A$ is not invertible, so that $A^T A\mathbf{x} = \mathbf{0}$ for some $\mathbf{x} \neq \mathbf{0}$. Multiply both sides with \mathbf{x}^T to get

$$\mathbf{x}^T A^T A\mathbf{x} = (A\mathbf{x})^T A\mathbf{x} = \|A\mathbf{x}\|^2 = 0,$$

which implies that $A\mathbf{x} = \mathbf{0}$. Since the columns of A are independent, this shows that $\mathbf{x} = \mathbf{0}$. A contradiction! \square

Example 52.

- (a) What is the matrix P for projecting onto $W = \text{span}\left\{\begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}\right\}$?
- (b) Using P , what is the orthogonal projection of $\begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$ onto W ?
- (c) Using P , what is the orthogonal projection of $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ onto W ?

Solution.

- (a) Note that $W = \text{col}(A)$ for $A = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}$ and that $A^T A = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$. Thus $(A^T A)^{-1} = \frac{1}{84} \begin{bmatrix} 5 & -1 \\ -1 & 17 \end{bmatrix}$.

$$P = A(A^T A)^{-1} A^T = \frac{1}{84} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 5 & -1 \\ -1 & 17 \end{bmatrix} \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} = \frac{1}{21} \begin{bmatrix} 20 & -2 & 4 \\ -2 & 17 & 8 \\ 4 & 8 & 5 \end{bmatrix}$$

- (b) The orthogonal projection of $\begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$ onto W is $P \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \frac{1}{21} \begin{bmatrix} 20 & -2 & 4 \\ -2 & 17 & 8 \\ 4 & 8 & 5 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \frac{1}{21} \begin{bmatrix} 84 \\ 84 \\ 63 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 3 \end{bmatrix}$.

Note. Of course, that agrees with what our computations in Example 49. Note that computing P is more work than what we did in Example 49. However, after having computed P once, we can easily project many vectors onto W .

- (c) The orthogonal projection of $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ onto W is $P \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \frac{1}{21} \begin{bmatrix} 20 & -2 & 4 \\ -2 & 17 & 8 \\ 4 & 8 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \frac{1}{21} \begin{bmatrix} 20 \\ -2 \\ 4 \end{bmatrix}$.

Check. The error $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \frac{1}{21} \begin{bmatrix} 20 \\ -2 \\ 4 \end{bmatrix} = \frac{1}{21} \begin{bmatrix} 1 \\ 2 \\ -4 \end{bmatrix}$ is indeed orthogonal to both $\begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$.

Example 53. (extra)

- (a) What is the matrix P for projecting onto $W = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}\right\}$?
- (b) Using the projection matrix, project $\begin{bmatrix} 2 \\ 3 \\ 3 \end{bmatrix}$ onto $W = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}\right\}$.

Solution.

(a) Choosing $A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix}$, the projection matrix P is $A(A^T A)^{-1} A^T = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix}$

$$= \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \frac{1}{8} \begin{bmatrix} 3 & -1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix} = \frac{1}{8} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & 2 \\ 2 & -4 & 2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{bmatrix}.$$

Comment. We can choose A in any way such that its columns are a basis for W . The final projection matrix will always be the same.

(b) The projection is $\frac{1}{2} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \\ 3 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 5 \\ 6 \\ 5 \end{bmatrix}$.

Check. The error $\begin{bmatrix} 2 \\ 3 \\ 3 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 5 \\ 6 \\ 5 \end{bmatrix} = \begin{bmatrix} -1/2 \\ 0 \\ 1/2 \end{bmatrix}$ is indeed orthogonal to W .

Example 54. If P is a projection matrix, then what is P^2 ?

For instance. For P as in Example 53, $P^2 = \frac{1}{4} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{bmatrix}^2 = \frac{1}{2} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{bmatrix} = P$.

Solution. Can you see why it is always true that $P^2 = P$?

[Recall that P projects a vector onto a space W (actually, $W = \text{col}(P)$). Hence P^2 takes a vector \mathbf{b} , projects it onto W to get $\hat{\mathbf{b}}$, and then projects $\hat{\mathbf{b}}$ onto W again. But the projection of $\hat{\mathbf{b}}$ onto W is just $\hat{\mathbf{b}}$ (why?!), so that P^2 always has the exact same effect as P . Therefore, $P^2 = P$.]

Projecting onto 1-dimensional spaces

When we project onto a 1-dimensional space $\text{span}\{\mathbf{w}\}$, we usually just say that we are projecting onto \mathbf{w} .

The (orthogonal) projection of \mathbf{v} onto \mathbf{w} is $\frac{\mathbf{w} \cdot \mathbf{v}}{\|\mathbf{w}\|^2} \mathbf{w}$.

Why? Replace \mathbf{b} with \mathbf{v} and A with \mathbf{w} in our general projection matrix formula to get $\mathbf{w}(\mathbf{w}^T \mathbf{w})^{-1} \mathbf{w}^T \mathbf{v}$, which equals $\frac{\mathbf{w} \cdot \mathbf{v}}{\|\mathbf{w}\|^2} \mathbf{w}$ (note that $\mathbf{w}^T \mathbf{v} = \mathbf{w} \cdot \mathbf{v}$ and $\mathbf{w}^T \mathbf{w} = \|\mathbf{w}\|^2$ are scalars).

Comment. If you have taken Calculus 3, you have seen that formula before. Most likely, you were deriving it using angles at that time. Namely, the dot product has the following connection to angles:

$$\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos \theta \text{ where } \theta \in [0, \pi] \text{ is the angle between } \mathbf{v} \text{ and } \mathbf{w}$$

Why? You can derive this by repeating what we did, right after Definition 23 to show that \mathbf{v} and \mathbf{w} are orthogonal if and only if $\mathbf{v} \cdot \mathbf{w} = 0$. Just replace Pythagoras with the law of cosines ($c^2 = a^2 + b^2 - 2ab \cos \theta$ holds in any triangle!).


Two obvious cases. Observe that the cases $\theta = 0$ and $\theta = 90^\circ$ are clearly true.

We will not discuss angles much further in this class. Just in case it is helpful, here is the typical argument given in Calculus 3 to determine the projection $\text{proj}_{\mathbf{w}} \mathbf{v}$ of \mathbf{v} onto \mathbf{w} :

From the sketch, we see that “error” = $v - \text{proj}_w v$ and that this error is orthogonal to w .

Basic trigonometry tells us that the length of $\text{proj}_w v$ is $\|v\| \cos\theta$. Hence:

$$\begin{aligned} \text{proj}_w v &= \underbrace{\|v\| \cos\theta}_{\text{length}} \underbrace{\frac{w}{\|w\|}}_{\text{direction}} \\ &= \frac{\|v\| \|w\| \cos\theta}{\|w\|} \frac{w}{\|w\|} = \left(\frac{v \cdot w}{\|w\|^2} \right) w \end{aligned}$$



Orthogonal bases

Review. Vectors v_1, \dots, v_n are a **basis** for V .

$\iff V = \text{span}\{v_1, \dots, v_n\}$ and v_1, \dots, v_n are linearly independent.

\iff Any vector w in V can be written as $w = c_1 v_1 + \dots + c_n v_n$ in a unique way.

The latter is the practical reason why we care so much about bases!

V could be some abstract vector space (of polynomials or Fourier series), meaning that vectors are abstract objects and not just our usual column vectors. However, as soon as we pick a basis of V , then we can represent every (abstract) vector w by the (usual) column vector $(c_1, c_2, \dots, c_n)^T$.

This means all of our results can be used, too, when working with these abstract spaces!

Definition 55. A basis v_1, \dots, v_n of a vector space V is an **orthogonal basis** if the vectors are (pairwise) orthogonal. If, in addition, the basis vectors have length 1, then this is called an **orthonormal basis**.

Example 56. The standard basis $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ is an orthonormal basis for \mathbb{R}^3 .

Example 57. Are the vectors $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ an orthogonal basis for \mathbb{R}^3 ? Is it orthonormal?

Solution. $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = 0, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0.$

So, this is an orthogonal basis.

On the other hand, the vectors do not all have length 1, so that this basis is not orthonormal.

Note. Orthogonal vectors are always linearly independent (see next class). Here, this certifies that the three vectors are linearly independent (and hence a basis for \mathbb{R}^3).

Normalize the vectors to produce an orthonormal basis.

Solution.

$$\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \text{ has length } \sqrt{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} = \sqrt{2} \implies \text{normalized: } \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \text{ has length } \sqrt{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} = \sqrt{2} \implies \text{normalized: } \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \text{ has length } \sqrt{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}} = 1 \implies \text{is already normalized: } \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

The resulting orthonormal basis is $\frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.

Review. The **projection matrix** for projecting onto $\text{col}(A)$ is $P = A(A^T A)^{-1} A^T$.

Example 58. True or false? If P is the matrix for projecting onto W , then $W = \text{col}(P)$.

Solution. True!

Why? The columns of P are the projections of the standard basis vectors and hence in W . On the other hand, for any vector w in W , we have $Pw = w$ so that w is a combination of the columns of P .

[This may take several readings to digest but do read (or ask) until it makes sense!]

In particular. $\text{rank}(P) = \dim W$ (because, for any matrix, $\text{rank}(A) = \dim \text{col}(A)$)

Theorem 59. Suppose that v_1, \dots, v_n are nonzero and pairwise orthogonal. Then v_1, \dots, v_n are linearly independent.

Proof. Suppose that $c_1 v_1 + \dots + c_n v_n = 0$. In order to show that v_1, \dots, v_n are independent, we need to show that $c_1 = c_2 = \dots = c_n = 0$.

Take the dot product of v_1 with both sides:

$$\begin{aligned} 0 &= v_1 \cdot (c_1 v_1 + \dots + c_n v_n) \\ &= c_1 v_1 \cdot v_1 + c_2 v_1 \cdot v_2 + \dots + c_n v_1 \cdot v_n \\ &= c_1 v_1 \cdot v_1 = c_1 \|v_1\|^2 \end{aligned}$$

But $\|v_1\| \neq 0$ and hence $c_1 = 0$. Likewise, we find $c_2 = 0, \dots, c_n = 0$. Hence, the vectors are independent. \square

Comment. Note that this result is intuitively obvious: if the vectors were linearly dependent, then one of them could be written as a linear combination of the others. However, all these other vectors (and hence any combination of them) are orthogonal to it.

Orthogonal projections if we have an orthogonal basis

Lemma 60. (orthogonal projection if we have an orthogonal basis)

If v_1, \dots, v_n are orthogonal, then the orthogonal projection of w onto $\text{span}\{v_1, \dots, v_n\}$ is

$$\hat{w} = \underbrace{\frac{w \cdot v_1}{v_1 \cdot v_1} v_1}_{\text{proj of } w \text{ onto } v_1} + \dots + \underbrace{\frac{w \cdot v_n}{v_n \cdot v_n} v_n}_{\text{proj of } w \text{ onto } v_n}$$

Proof. It suffices to show that the error $w - \hat{w}$ is orthogonal to each v_i . Indeed:

$$(w - \hat{w}) \cdot v_i = \left(w - \frac{w \cdot v_1}{v_1 \cdot v_1} v_1 - \dots - \frac{w \cdot v_n}{v_n \cdot v_n} v_n \right) \cdot v_i = w \cdot v_i - \frac{w \cdot v_i}{v_i \cdot v_i} v_i \cdot v_i = 0.$$

\square

Important consequence. If v_1, \dots, v_n is an orthogonal basis of V , and w is in V , then

$$w = c_1 v_1 + \dots + c_n v_n \quad \text{with} \quad c_j = \frac{w \cdot v_j}{v_j \cdot v_j}$$

If the v_1, \dots, v_n are a basis, but not orthogonal, then we have to solve a system of equations to find the c_i . That is a lot more work than simply computing a few dot products.

Note. In other words, w decomposes as the sum of its projections onto each basis vector.

Note. If v_1, \dots, v_n are orthonormal, then the denominators are all 1.

Example 61. What is the projection of $\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix}$ onto $W = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$ with $\mathbf{v}_1 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$?

Comment. We know how to do this using least squares. (Do it for practice!)

However, realizing that \mathbf{v}_1 and \mathbf{v}_2 are orthogonal makes things easier.

[Actually, here, it is obvious what the projection is going to be if we realized that W is the x - y -plane.]

Solution. (using orthogonality) Because \mathbf{v}_1 and \mathbf{v}_2 are orthogonal, the projection is

$$\underbrace{\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{\text{projection onto } \mathbf{v}_1} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \underbrace{\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}_{\text{projection onto } \mathbf{v}_2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \frac{-4}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{10}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 7 \\ 0 \end{bmatrix}.$$

Important note. Note that, at this point, we can easily extend $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ to an orthogonal basis of \mathbb{R}^3 :

That is because the error $\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix} - \begin{bmatrix} 3 \\ 7 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 4 \end{bmatrix}$ is orthogonal to both of the existing basis vectors.

Therefore $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 4 \end{bmatrix}$ is an orthogonal basis of \mathbb{R}^3 .

This observation underlies the Gram-Schmidt process, which we will discuss next class.

Example 62. Express $\underbrace{\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix}}_{\mathbf{x}}$ in terms of the basis $\underbrace{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{\mathbf{v}_1}, \underbrace{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}_{\mathbf{v}_2}, \underbrace{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}_{\mathbf{v}_3}$.

Solution. Because $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ is an orthogonal basis of \mathbb{R}^3 , we get (much as in the previous example):

$$\begin{aligned} \begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix} &= c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\ &= \underbrace{\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{\text{projection of } \mathbf{x} \text{ onto } \mathbf{v}_1} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \underbrace{\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}_{\text{projection of } \mathbf{x} \text{ onto } \mathbf{v}_2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \underbrace{\begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}_{\text{projection of } \mathbf{x} \text{ onto } \mathbf{v}_3} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\ &= \frac{-4}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{10}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \frac{4}{1} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

Because we spelled out all the details this looks more involved than it is. We only computed 6 dot products!

Alternative. We could have solved $\begin{bmatrix} 1 & 1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 7 \\ 4 \end{bmatrix}$ to also find $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} -2 \\ 5 \\ 4 \end{bmatrix}$.

The numbers are particularly easy here but in general, to find this solution, we have to go through the entire process of Gaussian elimination. On the other hand, if we have an orthogonal basis, the former approach requires less work, because it is just computing a few dot products.

Example 63. Express $\begin{bmatrix} 2 \\ 5 \\ 1 \end{bmatrix}$ in terms of the basis $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$.

Solution. This is not an orthogonal basis, so we cannot proceed as in the previous example.

To write $\begin{bmatrix} 2 \\ 5 \\ 1 \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} + c_3 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, we need to solve $\begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 5 \\ 1 \end{bmatrix}$.

Solving that system (do it!), we find $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}$.

Review. If v_1, \dots, v_n are orthogonal, the orthogonal projection of w onto $\text{span}\{v_1, \dots, v_n\}$ is

$$\hat{w} = \frac{w \cdot v_1}{v_1 \cdot v_1} v_1 + \dots + \frac{w \cdot v_n}{v_n \cdot v_n} v_n.$$

Example 64.

(a) Project $\begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ onto $W = \text{span}\left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} \right\}$.

(b) Express $\begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ in terms of the basis $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ -5 \end{bmatrix}$.

Solution.

(a) The projection is $\frac{8}{6} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + \frac{4}{5} \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}$. (Each coefficient is obtained as the quotient of two dot products.)

(b) $\begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} = \frac{8}{6} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} + \frac{4}{5} \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} + \frac{5}{30} \begin{bmatrix} 1 \\ 2 \\ -5 \end{bmatrix}$

Gram–Schmidt

(Gram–Schmidt orthogonalization)

Given a basis w_1, w_2, \dots for W , we produce an orthogonal basis q_1, q_2, \dots for W as follows:

- $q_1 = w_1$
- $q_2 = w_2 - \left(\begin{smallmatrix} \text{projection of} \\ w_2 \text{ onto } q_1 \end{smallmatrix} \right)$
- $q_3 = w_3 - \left(\begin{smallmatrix} \text{projection of} \\ w_3 \text{ onto } q_1 \end{smallmatrix} \right) - \left(\begin{smallmatrix} \text{projection of} \\ w_3 \text{ onto } q_2 \end{smallmatrix} \right)$
- $q_4 = \dots$

Note. Since q_1, q_2 are orthogonal, $\left(\begin{smallmatrix} \text{projection of} \\ w_3 \text{ onto } \text{span}\{q_1, q_2\} \end{smallmatrix} \right) = \left(\begin{smallmatrix} \text{projection of} \\ w_3 \text{ onto } q_1 \end{smallmatrix} \right) + \left(\begin{smallmatrix} \text{projection of} \\ w_3 \text{ onto } q_2 \end{smallmatrix} \right)$.

Important comment. When working numerically on a computer it actually saves time to compute an orthonormal basis q_1, q_2, \dots by the same approach but always normalizing each q_i along the way. The reason this saves time is that now the projections onto q_i only require a single dot product (instead of two). This is called **Gram–Schmidt orthonormalization**. When working by hand, it is usually simpler to wait until the end to normalize (so as to avoid).

Note. When normalizing, the orthonormal basis q_1, q_2, \dots is the unique one (up to \pm signs) with the property that $\text{span}\{q_1, q_2, \dots, q_k\} = \text{span}\{w_1, w_2, \dots, w_k\}$ for all $k = 1, 2, \dots$

Example 65. Using Gram–Schmidt, find an orthogonal basis for $W = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}\right\}$.

Solution. We already have the basis $w_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$, $w_2 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$ for W . However, that basis is not orthogonal.

We can construct an orthogonal basis q_1, q_2 for W as follows:

- $q_1 = w_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$
- $q_2 = w_2 - \left(\text{projection of } w_2 \text{ onto } q_1\right) = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} - \frac{1}{3}\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \frac{1}{3}\begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix}$

Note. q_2 is the error of the projection of w_2 onto q_1 . This guarantees that it is orthogonal to q_1 .
On the other hand, since q_2 is a combination of w_2 and q_1 , we know that q_2 actually is in W .

We have thus found the orthogonal basis $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \frac{2}{3}\begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$ for W (if we like, we can, of course, drop that $\frac{2}{3}$).

Important comment. By normalizing, we get an orthonormal basis for W : $\frac{1}{\sqrt{3}}\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \frac{1}{\sqrt{6}}\begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$.

Practical comment. When implementing Gram–Schmidt on a computer, it is beneficial (slightly less work) to normalize each q_i during the Gram–Schmidt process. This typically introduces square roots, which is why normalizing at the end is usually preferable when working by hand.

Comment. There are, of course, many orthogonal bases q_1, q_2 for W . Up to the length of the vectors, ours is the unique one with the property that $\text{span}\{q_1\} = \text{span}\{w_1\}$ and $\text{span}\{q_1, q_2\} = \text{span}\{w_1, w_2\}$.

A matrix Q has orthonormal columns $\iff Q^T Q = I$

Why? Let q_1, q_2, \dots be the columns of Q . By the way matrix multiplication works, the entries of $Q^T Q$ are dot products of these columns:

$$\begin{bmatrix} - & q_1^T & - \\ - & q_2^T & - \\ \vdots & & \end{bmatrix} \begin{bmatrix} | & | & \dots \\ q_1 & q_2 & \dots \\ | & | & \dots \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \ddots \end{bmatrix}$$

Hence, $Q^T Q = I$ if and only if the dot products $q_i^T q_j = 0$ (that is, the columns are orthogonal), for $i \neq j$, and $q_i^T q_i = 1$ (that is, the columns are normalized).

Example 66. $Q = \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{6} \\ 1/\sqrt{3} & -2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{6} \end{bmatrix}$ obtained from Example 65 satisfies $Q^T Q = I$.

The QR decomposition

Just like the LU decomposition encodes the steps of Gaussian elimination, the QR decomposition encodes the steps of Gram–Schmidt.

(QR decomposition) Every $m \times n$ matrix A of rank n can be decomposed as $A = QR$, where

- Q has orthonormal columns, $(m \times n)$
- R is upper triangular and invertible. $(n \times n)$

How to find Q and R ?

- Gram–Schmidt orthonormalization on (columns of) A , to get (columns of) Q
- $R = Q^T A$

Why? If $A = QR$, then $Q^T A = Q^T QR$ which simplifies to $R = Q^T A$ (since $Q^T Q = I$).

The decomposition $A = QR$ is unique if we require the diagonal entries of R to be positive (and this is exactly what happens when applying Gram–Schmidt).

Practical comment. Actually, no extra work is needed for computing R . All of its entries have been computed during Gram–Schmidt.

Variations. We can also arrange things so that Q is an $m \times m$ orthogonal matrix and R a $m \times n$ upper triangular matrix. This is a tiny bit more work (and not required for many applications): we need to complement “our” Q with additional orthonormal columns and add corresponding zero rows to R . For square matrices this makes no difference.

Example 67. Determine the QR decomposition of $A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix}$.

Solution. The first step is Gram–Schmidt orthonormalization on the columns of A . We then use the resulting orthonormal vectors (they need to be normalized!) as the columns of Q .

We already did Gram–Schmidt in Example 65: from that work, we have $Q = \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{6} \\ 1/\sqrt{3} & -2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{6} \end{bmatrix}$.

Hence, $R = Q^T A = \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \\ 1/\sqrt{6} & -2/\sqrt{6} & 1/\sqrt{6} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 3/\sqrt{3} & 1/\sqrt{3} \\ 0 & 4/\sqrt{6} \end{bmatrix}$.

Comment. As commented earlier, the entries of R have actually all been computed during Gram–Schmidt, so that, if we pay attention, we could immediately write down R (no extra work required). Looking back at Example 65, can you see this?

Check. Indeed, $QR = \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{6} \\ 1/\sqrt{3} & -2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{6} \end{bmatrix} \begin{bmatrix} 3/\sqrt{3} & 1/\sqrt{3} \\ 0 & 4/\sqrt{6} \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 1 \end{bmatrix}$ equals A .

Example 68. Using Gram–Schmidt, find an orthogonal basis for $W = \text{span} \left\{ \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right\}$.

Solution. We begin with the (not orthogonal) basis $w_1 = \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \end{bmatrix}$, $w_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}$, $w_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$.

We then construct an orthogonal basis q_1, q_2, q_3 :

- $q_1 = w_1 = \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \end{bmatrix}$
- $q_2 = w_2 - \left(\begin{matrix} \text{projection of} \\ w_2 \text{ onto } q_1 \end{matrix} \right) = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \frac{3}{9} \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 0 \\ 0 \end{bmatrix}$
- $q_3 = w_3 - \left(\begin{matrix} \text{projection of } w_3 \\ \text{onto } \text{span}\{q_1, q_2\} \right) = w_3 - \left(\begin{matrix} \text{projection of} \\ w_3 \text{ onto } q_1 \end{matrix} \right) - \left(\begin{matrix} \text{projection of} \\ w_3 \text{ onto } q_2 \end{matrix} \right)$
 $= \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{3}{9} \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \end{bmatrix} - \frac{2}{4} \begin{bmatrix} 2 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$

Make sure you understand how q_3 was designed to be orthogonal to both q_1 and q_2 !

Also note that breaking up the projection onto $\text{span}\{q_1, q_2\}$ into the projections onto q_1 and q_2 is only possible because q_1 and q_2 are orthogonal.

Hence, $\left\{ \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right\}$ is an orthogonal basis of W .

Important. Normalizing, we obtain an orthonormal basis: $\left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} \right\}$.

Example 69. Determine the QR decomposition of $A = \begin{bmatrix} 0 & 2 & 1 \\ 3 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$.

Solution. The first step is Gram–Schmidt orthonormalization on the columns of A . We then use the resulting orthonormal vectors as the columns of Q .

We already did Gram–Schmidt in Example 68: from that work, we have $Q = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1/\sqrt{2} \\ 0 & 0 & 1/\sqrt{2} \end{bmatrix}$.

Hence, $R = Q^T A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 0 & 2 & 1 \\ 3 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & \sqrt{2} \end{bmatrix}$.

Comment. As commented earlier, the entries of R have actually all been computed during Gram–Schmidt, so that, if we pay attention, we could immediately write down R (no extra work required). Looking back at Example 68, can you see this?

Letting Sage do the work for us.

```
Sage] A = matrix(QQbar, [[0,2,1],[3,1,1],[0,0,1],[0,0,1]])
```

```
Sage] A.QR(full=false)
```

$$\left(\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0.7071067811865475? \\ 0 & 0 & 0.7071067811865475? \end{pmatrix}, \begin{pmatrix} 3 & 1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 1.414213562373095? \end{pmatrix} \right)$$

Comment. Can you figure out what happens if you omit the `full=false`? Check out the comment under **Variations** for the QR decomposition in the previous lecture sketch. On the other hand, the `QQbar` is telling Sage to compute with algebraic numbers (instead of just rational numbers); if omitted, it would complain that square roots are not available

Example 70. (extra) Determine the QR decomposition of $A = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & -5 \\ 0 & 3 & 6 \end{bmatrix}$.

Solution. We first apply Gram–Schmidt orthonormalization to the columns of A . For a variation, like a computer, we normalize after each step (rather than normalize at the end):

- $\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, so that $\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$.
- $\mathbf{b}_2 = \begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix} - \left(\begin{bmatrix} 2 \\ 0 \\ 3 \end{bmatrix} \cdot \mathbf{q}_1 \right) \mathbf{q}_1 = \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$, so that $\mathbf{q}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.
- $\mathbf{b}_3 = \begin{bmatrix} 4 \\ -5 \\ 6 \end{bmatrix} - \left(\begin{bmatrix} 4 \\ -5 \\ 6 \end{bmatrix} \cdot \mathbf{q}_1 \right) \mathbf{q}_1 - \left(\begin{bmatrix} 4 \\ -5 \\ 6 \end{bmatrix} \cdot \mathbf{q}_2 \right) \mathbf{q}_2 = \begin{bmatrix} 0 \\ -5 \\ 0 \end{bmatrix}$, so that $\mathbf{q}_3 = \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$.

Therefore, $Q = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}$. Finally, $R = Q^T A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & -5 \\ 0 & 3 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{bmatrix}$.

In conclusion, we have found the QR decomposition: $\underbrace{\begin{bmatrix} 1 & 2 & 4 \\ 0 & 0 & -5 \\ 0 & 3 & 6 \end{bmatrix}}_A = \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}}_Q \underbrace{\begin{bmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{bmatrix}}_R$

Comment. As noted before, we actually could write down R without any additional computation. Indeed, realize that the second column of R , that is $[2, 3, 0]^T$ means that

$$\text{2nd col of } A = 2\mathbf{q}_1 + 3\mathbf{q}_2.$$

Which we already knew from our computation of \mathbf{q}_2 ! Also, by construction, we know that the second column of A is a linear combination of \mathbf{q}_1 and \mathbf{q}_2 only, and that \mathbf{q}_3 enters the story later on. This corresponds to the fact that R is always upper triangular.

Letting Sage do the work for us.

```
Sage] A = matrix(QQbar, [[1,2,4], [0,0,-5], [0,3,6]])
```

```
Sage] A.QR()
```

$$\left(\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{pmatrix} \right)$$

Comment. The `QQbar` is telling Sage to compute with algebraic numbers (instead of just rational numbers); in general, if omitted, it would complain that square roots are not available (because the matrices Q and R typically involve square roots). Here, we are lucky that square roots didn't creep in.

Example 71. (extra) Find the QR decomposition of $A = \begin{bmatrix} 1 & 1 & 2 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$.

Solution. (final answer only) $A = QR$ with $Q = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \end{bmatrix}$ and $R = \begin{bmatrix} \sqrt{2} & \frac{1}{\sqrt{2}} & \sqrt{2} \\ 0 & \frac{1}{\sqrt{2}} & \sqrt{2} \\ 0 & 0 & 1 \end{bmatrix}$.

Example 72. One practical application of the QR decomposition is solving systems of linear equations.

$$\begin{aligned} Ax = b &\iff QRx = b && \text{(now, multiply with } Q^T \text{ from the left)} \\ &\implies Rx = Q^T b \end{aligned}$$

The last system is triangular and can be solved by back substitution.

A couple of comments are in order:

- If A is $n \times n$ and invertible, then the " \implies " is actually a " \iff ".
- The equation $Rx = Q^T b$ is always consistent! (Recall that R is invertible.)
Indeed, if A is not $n \times n$ or not invertible, then $Rx = Q^T b$ gives the least squares solutions!
Why? $A^T A \hat{x} = A^T b \iff \underbrace{(QR)^T Q R \hat{x}}_{=R^T Q^T Q R} = (QR)^T b \iff R^T R \hat{x} = R^T Q^T b \iff R \hat{x} = Q^T b$
[For the last step we need that R is invertible, which is always the case when A is $m \times n$ of rank n .]

- So, how does the QR way of solving linear systems compare to our beloved Gaussian elimination (LU)?
It turns out that QR is a little slower than LU but makes up for it in "numerical stability".

What does that mean? When computing numerically, we use floating point arithmetic and approximate each number by an expression of the form $0.1234 \cdot 10^{-16}$. A certain (fixed) number of bits is used to store the part 0.1234 (here, 4 decimal places of accuracy) as well as the exponent -16 .

Now, here is something terrible that can happen in numerical computations: mathematically, the quantities x and $(x+1) - 1$ are exactly the same. However, numerically, they might not. Take, for instance, $x = 0.1234 \cdot 10^{-6}$. Then, to an accuracy of 4 decimal places, $x+1 = 0.1000 \cdot 10^1$, so that $(x+1) - 1 = 0.0000$. But $x \neq 0$. We completely lost all the information about x .

To be numerically stable, an algorithm must avoid issues like that.

\hat{x} is a least squares solution of $Ax = b$
 $\iff R\hat{x} = Q^T b$ (where $A = QR$)

Review. A matrix A has orthonormal columns $\iff A^T A = I$.

Example 73. Suppose Q has orthonormal columns. What is the projection matrix P for orthogonally projecting onto $\text{col}(Q)$?

Solution. Recall that, to project onto $\text{col}(A)$, the projection matrix is $P = A(A^T A)^{-1} A^T$.

Since $Q^T Q = I$, to project onto $\text{col}(Q)$, the projection matrix is $P = Q Q^T$.

Comment. A familiar special case is when we project onto a unit vector q : in that case, the projection of b onto q is $(q \cdot b)q = q(q^T b) = (qq^T)b$, so the projection matrix here is qq^T .

Comment. In particular, if Q is not square, then $Q^T Q = I$ but $Q Q^T \neq I$. In some sense, $Q Q^T$ still “tries” to be as close to the identity as possible: since it is the matrix projecting onto $\text{col}(Q)$ it does act like the identity for vectors in $\text{col}(Q)$. (Vectors not in $\text{col}(Q)$ are sent to their projection, that is, the closest to themselves while restricted to $\text{col}(Q)$.)

Example 74. Suppose A is invertible. What is the projection matrix P for orthogonally projecting onto $\text{col}(A)$?

Solution. If A is an invertible $n \times n$ matrix, then $\text{col}(A) = \mathbb{R}^n$ (because the n columns of A are linearly independent and hence form a basis for \mathbb{R}^n).

Since $\text{col}(A)$ is the entire space we are not really projecting at all: every vector is sent to itself.

In particular, the projection matrix is $P = I$.

Definition 75. An **orthogonal matrix** is a square matrix with orthonormal columns.

[This is not a typo (but a confusing convention): the columns need to be orthonormal, not just orthogonal.]

An $n \times n$ matrix Q is orthogonal $\iff Q^T Q = I$

In other words, $Q^{-1} = Q^T$.

Example 76. What can we say about $\det(Q)$ if Q is orthogonal?

Solution. Write $d = \det(Q)$. Since $Q^{-1} = Q^T$, we have $\frac{1}{d} = d$ (recall that $\det(Q^{-1}) = 1 / \det(Q)$ and $\det(Q^T) = \det(Q)$) or, equivalently, $d^2 = 1$. Hence, $d = \pm 1$.

Both of these are possible as the examples $Q = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $Q = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ illustrate.

Review: More on diagonalization

Example 77. (review) If A is a 2×2 matrix with $\det(A) = -8$ and eigenvalue 4. What is the second eigenvalue?

Solution. Recall that $\det(A)$ is the product of the eigenvalues (see below). Hence, the second eigenvalue is -2 .

$\det(A)$ is the product of the eigenvalues of A .

Why? Recall how we determine the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ of an $n \times n$ matrix A . We compute the characteristic polynomial $\det(A - \lambda I)$ and determine the λ_i as the roots of that polynomial.

That means that we have the factorization $\det(A - \lambda I) = (\lambda_1 - \lambda)(\lambda_2 - \lambda) \cdots (\lambda_n - \lambda)$. Now, set $\lambda = 0$ to conclude that $\det(A) = \lambda_1 \lambda_2 \cdots \lambda_n$.

Example 78.

- (a) Determine the eigenspaces of the symmetric matrix $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$.
- (b) Diagonalize A as $A = PDP^T$.

Solution.

- (a) The characteristic polynomial is $\begin{vmatrix} 1-\lambda & 3 \\ 3 & 1-\lambda \end{vmatrix} = (\lambda-4)(\lambda+2)$, and so A has eigenvalues $4, -2$.

The 4 -eigenspace is $\text{null}\left(\begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix}\right)$ has basis $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

The -2 -eigenspace is $\text{null}\left(\begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix}\right)$ has basis $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$.

Important observation. The 4 -eigenvector $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and the -2 -eigenvector $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ are orthogonal!

Review. The product of all eigenvalues $-2 \cdot 4 = -8$ equals the determinant $\det(A) = 1 - 9 = -8$.

- (b) Note that a usual diagonalization is of the form $A = PDP^{-1}$.
We need to choose P so that $P^{-1} = P^T$, which means that P must be **orthogonal** (meaning orthonormal columns). [Choosing such a P is only possible if the eigenspaces of A are orthogonal.]

Hence, we normalize the two eigenvectors to $\frac{1}{\sqrt{2}}\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\frac{1}{\sqrt{2}}\begin{bmatrix} -1 \\ 1 \end{bmatrix}$.

With $P = \frac{1}{\sqrt{2}}\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$ and $D = \begin{bmatrix} 4 & 0 \\ 0 & -2 \end{bmatrix}$, we then have $A = PDP^T$.

Example 79. (homework) Diagonalize, if possible, the matrices

(a) $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, and

(b) $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$.

We will briefly discuss these next class.

Example 80. (review) In Example 15, we diagonalized $A = \begin{bmatrix} 4 & 0 & 2 \\ 2 & 2 & 2 \\ 1 & 0 & 3 \end{bmatrix}$ as $A = PDP^{-1}$.

We found that one choice for P and D is $P = \begin{bmatrix} 2 & 0 & -1 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$, $D = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

Spell out what that tells us about A !

Solution. The diagonal entries 5, 2, 2 of D are the eigenvalues of A .

The columns of P are corresponding eigenvectors of A .

- $\begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$ is a 5-eigenvector of A (that is, $A \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} = 5 \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$).
- The 2-eigenspace of A is 2-dimensional. A basis is $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$.

Lemma 81. A matrix A is diagonalizable if and only if, for every eigenvalue λ that is k times repeated, the λ -eigenspace of A has dimension k .

In short, an $n \times n$ matrix A is diagonalizable if and only if there exists a basis of \mathbb{R}^n consisting of eigenvectors of A (i.e. “there are enough eigenvectors”).

The next two examples illustrate that not all matrices are diagonalizable and that, even if a real matrix is diagonalizable, the eigenvalues and eigenvectors might be complex.

Example 82. What are the eigenvalues and eigenvectors of $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$? Is A diagonalizable?

Solution. The characteristic polynomial is $\det\left(\begin{bmatrix} -\lambda & 1 \\ 0 & -\lambda \end{bmatrix}\right) = \lambda^2$, which has $\lambda = 0$ as a double root.

However, the 0-eigenspace $\text{null}(A) = \text{span}\left\{\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right\}$ is only 1-dimensional.

As a consequence, A is not diagonalizable.

Example 83. What are the eigenvalues and eigenvectors of $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$? Is A diagonalizable?

Solution. The characteristic polynomial is $\det\left(\begin{bmatrix} -\lambda & -1 \\ 1 & -\lambda \end{bmatrix}\right) = \lambda^2 + 1 = (\lambda - i)(\lambda + i)$.

Hence, the eigenvalues are $\pm i$.

The i -eigenspace $\text{null}\left(\begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix}\right)$ has basis $\begin{bmatrix} i \\ 1 \end{bmatrix}$.

The $-i$ -eigenspace $\text{null}\left(\begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix}\right)$ has basis $\begin{bmatrix} -i \\ 1 \end{bmatrix}$.

Thus, A has the diagonalization $A = PDP^{-1}$ with $D = \begin{bmatrix} i & \\ & -i \end{bmatrix}$ and $P = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix}$.

The spectral theorem

Recall that a matrix A is symmetric if and only if $A^T = A$.

Theorem 84. (spectral theorem, long version) Suppose A is a symmetric matrix.

- A is always diagonalizable.
- All eigenvalues of A are real.
- The eigenspaces of A are orthogonal.

Comment. The eigenspaces of A being orthogonal means that eigenvectors for different eigenvalues are always orthogonal.

Important consequence. In the diagonalization $A = PDP^{-1}$, we can choose P to be orthogonal (in which case $P^{-1} = P^T$). In that case, the diagonalization takes the special form $A = PDP^T$, where P is orthogonal and D is diagonal.

Example 85. (again) Diagonalize the symmetric matrix $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ as $A = PDP^T$.

Solution. See Example 78 for a solution that illustrates how to diagonalize any symmetric matrix.

Here, let us observe that (because the row sums are equal!) $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is a 4-eigenvector.

Because the eigenspaces are orthogonal (since A is symmetric!), $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ must be an eigenvector. $\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix}$, so the corresponding eigenvalue is -2 .

We normalize the two eigenvectors and use them as the columns of P , so that $P = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$ is an orthogonal matrix ($P^{-1} = P^T$). With $D = \begin{bmatrix} 4 & 0 \\ 0 & -2 \end{bmatrix}$ we then have $A = PDP^T$.

Example 86. Let A be a symmetric 2×2 matrix with 7-eigenvector $\begin{bmatrix} 2 \\ 5 \end{bmatrix}$ and $\det(A) = -21$. Determine the second eigenvalue and a corresponding eigenvector.

Solution. A has $-\frac{21}{7} = -3$ -eigenvector $\begin{bmatrix} -5 \\ 2 \end{bmatrix}$.

Comment. Recall that, because A is symmetric, the eigenvector must be orthogonal to $\begin{bmatrix} 2 \\ 5 \end{bmatrix}$.

[In general, $\begin{bmatrix} a \\ b \end{bmatrix}$ and $\begin{bmatrix} -b \\ a \end{bmatrix}$ are orthogonal.]

(spectral theorem, compact version) A symmetric matrix A can always be diagonalized as $A = PDP^T$, where P is orthogonal and D is diagonal (and both are real).

How? We proceed as in the diagonalization $A = PDP^{-1}$. We then arrange P to be orthogonal, by normalizing its columns. If there is a repeated eigenvalue, then we also need to make sure to pick an orthonormal basis for the corresponding eigenspace (for instance, using Gram–Schmidt).

Advanced comment. A matrix such that $A^T A = A A^T$ is called **normal**. For normal matrices, the (complex!) eigenspaces are again orthogonal to each other. However, normal matrices which are not symmetric will always have complex eigenvalues. (In that case, the orthogonal matrix P gets replaced with a unitary matrix, the complex version of orthogonal matrices, and the P^T becomes the conjugate transpose $P^* = \bar{P}^T$.)