

Example 110. Consider the sequence a_n defined by $a_{n+2} = 2a_{n+1} + 5a_n$ and $a_0 = 0, a_1 = 1$.

- (a) Determine the first few terms of the sequence.
- (b) Find a Binet-like formula for a_n .
- (c) Determine $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$.

Solution.

(a) 0, 1, 2, 9, 28, 101, 342, 1189, 4088, ...

(b) The recursion can be translated to $\begin{bmatrix} a_{n+2} \\ a_{n+1} \end{bmatrix} = \begin{bmatrix} 2 & 5 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a_{n+1} \\ a_n \end{bmatrix}$.

The eigenvalues of $\begin{bmatrix} 2 & 5 \\ 1 & 0 \end{bmatrix}$ are $1 \pm \sqrt{6}$.

Hence, $a_n = C_1(1 + \sqrt{6})^n + C_2(1 - \sqrt{6})^n$ and we only need to figure out the values of C_1 and C_2 .

Using the two initial conditions, we get two equations:

$$(a_0 = 0) \quad C_1 + C_2 = 0, \quad (a_1 = 1) \quad C_1(1 + \sqrt{6}) + C_2(1 - \sqrt{6}) = 1.$$

Solving, we find $C_1 = \frac{1}{2\sqrt{6}}$ and $C_2 = -\frac{1}{2\sqrt{6}}$ so that, in conclusion, $a_n = \frac{(1 + \sqrt{6})^n - (1 - \sqrt{6})^n}{2\sqrt{6}}$.

Comment. Alternatively, we could have proceeded as we did previously in the case of the Fibonacci numbers: starting with the recursion matrix M , we compute its diagonalization $M = PDP^{-1}$. Multiplying out $PD^nP^{-1} \begin{bmatrix} a_1 \\ a_0 \end{bmatrix}$, we obtain the Binet-like formula for a_n . However, this is more work than what we did.

(c) It follows from the Binet-like formula that $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 1 + \sqrt{6} \approx 3.44949$.

Comment. Actually, we don't need the Binet-like formula for this conclusion. Just the eigenvalues and the observation that C_1 cannot be 0 are enough. [We cannot have $C_1 = 0$, because then $a_n = C_2(1 - \sqrt{6})^n$ so that $a_0 = 0$ would imply $C_2 = 0$.]

Another brief look at projections (and reflections)

(projections) Suppose that M is the projection matrix for projecting onto a subspace W .

- The 1-eigenspace of M is W .
- The 0-eigenspace of M is W^\perp .

In particular, M is symmetric.

Why? By definition, the 1-eigenspace of M consists of those vectors that get projected to themselves. But those are precisely the vectors in W (recall that projecting a vector v onto W means producing the vector in W that is closest to v). Can you likewise spell out the situation for the 0-eigenspace?

Note that M cannot have further eigenvalues (because the dimensions of W and W^\perp already add up to the dimension of the space that we are working in).

Because the eigenvalues of M are real and the eigenspaces are orthogonal, the matrix M has a diagonalization of the form $M = PDP^T$ (make sure you can explain why!) which implies that M is symmetric (that's because $M^T = (PDP^T)^T = (P^T)^T D^T P^T = PDP^T = M$).

Example 111. Let A be the matrix for orthogonally projecting onto $W = \text{span}\left\{\begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}\right\}$.

- (a) Diagonalize A (without first computing A) as $A = PDP^{-1}$.
 (b) Diagonalize A as $A = PDP^T$.

Comment. This gives us yet another way to compute projection matrices: we can directly write down the matrices P, D for the diagonalization $A = PDP^T$. The main point here is that the diagonalization of a A nicely reveals all the information about the projection.

Solution.

- (a) The eigenvalues of A are $1, 1, 0$. The 1 -eigenspace of A is W (2-dimensional), and the 0 -eigenspace is W^\perp (1-dimensional).

We already have a basis for W . On the other hand, $W^\perp = \text{null}\left(\begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix}\right)$ has basis $\begin{bmatrix} -1/4 \\ -1/2 \\ 1 \end{bmatrix}$.

We therefore choose $D = \begin{bmatrix} 1 & & \\ & 1 & \\ & & 0 \end{bmatrix}$ and $P = \begin{bmatrix} 4 & 0 & -1/4 \\ 0 & 2 & -1/2 \\ 1 & 1 & 1 \end{bmatrix}$.

- (b) In order to achieve a diagonalization PDP^T we need to choose P to be orthogonal (which we can do here because the eigenspaces are orthogonal).

Applying Gram–Schmidt to the basis $w_1 = \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}$, $w_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$ (of the 1 -eigenspace), we construct the

orthogonal basis $q_1 = w_1 = \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}$, $q_2 = w_2 - \frac{w_2 \cdot q_1}{q_1 \cdot q_1} q_1 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix} - \frac{1}{17} \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix} = \frac{2}{17} \begin{bmatrix} -2 \\ 17 \\ 8 \end{bmatrix}$.

Next, we normalize the vectors $\begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}$, $\frac{1}{17} \begin{bmatrix} -4 \\ 34 \\ 16 \end{bmatrix}$, $\begin{bmatrix} -1/4 \\ -1/2 \\ 1 \end{bmatrix}$ to $\frac{1}{\sqrt{17}} \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}$, $\frac{1}{\sqrt{357}} \begin{bmatrix} -2 \\ 17 \\ 8 \end{bmatrix}$, $\frac{1}{\sqrt{21}} \begin{bmatrix} -1 \\ -2 \\ 4 \end{bmatrix}$.

We therefore choose $D = \begin{bmatrix} 1 & & \\ & 1 & \\ & & 0 \end{bmatrix}$ and $P = \begin{bmatrix} 4/\sqrt{17} & -2/\sqrt{357} & -1/\sqrt{21} \\ 0 & 17/\sqrt{357} & -2/\sqrt{21} \\ 1/\sqrt{17} & 8/\sqrt{357} & 4/\sqrt{21} \end{bmatrix}$.

By the way. Multiplying out $A = PDP^T$, we can find that $A = \frac{1}{21} \begin{bmatrix} 20 & -2 & 4 \\ -2 & 17 & 8 \\ 4 & 8 & 5 \end{bmatrix}$ as in Example 53.

Example 112. Let A be the matrix for orthogonally projecting onto $W = \text{span}\left\{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}\right\}$.

- (a) Diagonalize A (without first computing A) as $A = PDP^T$.
 (b) Is A invertible, orthogonal, symmetric?

Solution.

- (a) The eigenvalues of A are $1, 1, 0$. The 1 -eigenspace of A is W (2-dimensional), and the 0 -eigenspace is W^\perp (1-dimensional). Note that we are lucky and already have an orthogonal basis for W . On the other hand, $W^\perp = \text{null}\left(\begin{bmatrix} 1 & 1 & 1 \\ -1 & 0 & 1 \end{bmatrix}\right)$ has basis $\begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$.

We therefore choose $D = \begin{bmatrix} 1 & & \\ & 1 & \\ & & 0 \end{bmatrix}$ and, after normalizing columns, $P = \begin{bmatrix} 1/\sqrt{3} & -1/\sqrt{2} & 1/\sqrt{6} \\ 1/\sqrt{3} & 0 & -2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{2} & 1/\sqrt{6} \end{bmatrix}$.

- (b) A is not invertible (because 0 is an eigenvalue) and therefore also cannot be orthogonal. Like any projection matrix, A is symmetric.

By the way. Multiplying out $A = PDP^T$, we can find that $A = \frac{1}{6} \begin{bmatrix} 1 & -2 & 1 \\ -2 & 4 & -2 \\ 1 & -2 & 1 \end{bmatrix}$.