

# An application of modular forms to short random walks

AMS Fall Western Sectional Meeting — Tucson, Arizona

---

**Armin Straub**

October 28, 2012

University of Illinois  
at Urbana-Champaign

---

**Based on joint work with:**



Jon Borwein



James Wan



Wadim Zudilin

University of Newcastle, Australia

- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction

Q What is the distance traveled in  $n$  steps?

$p_n(x)$  probability density


$W_n(s)$   $s$ th moment

- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction


Q What is the distance traveled in  $n$  steps?

$p_n(x)$  probability density

$W_n(s)$   $s$ th moment



- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction




Q What is the distance traveled in  $n$  steps?

$p_n(x)$  probability density

$W_n(s)$   $s$ th moment

- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction




Q What is the distance traveled in  $n$  steps?

$p_n(x)$  probability density

$W_n(s)$   $s$ th moment

- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction




Q What is the distance traveled in  $n$  steps?

$p_n(x)$  probability density

$W_n(s)$   $s$ th moment

- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction




Q What is the distance traveled in  $n$  steps?

$p_n(x)$  probability density

$W_n(s)$   $s$ th moment

- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction




Q What is the distance traveled in  $n$  steps?

$p_n(x)$  probability density

$W_n(s)$   $s$ th moment

- $n$ -step uniform planar random walk in the plane:
  - $n$  steps, each of length 1,
  - taken in randomly chosen direction



**Q** What is the distance traveled in  $n$  steps?


$p_n(x)$  probability density  
 $W_n(s)$   $s$ th moment

**EG**



$$W_2(1) = \frac{4}{\pi}$$

- Karl Pearson asked for  $p_n(x)$  in Nature in 1905. This famous question coined the term **random walk**.
- **Asymptotic** answer by Lord Rayleigh:

$$p_n(x) \approx \frac{2x}{n} e^{-x^2/n}$$




EG  
p200




# Densities of short walks


$p_2$




$p_3$




$p_4$




$p_5$



$p_6$




$p_7$




# Densities of short walks


$p_2$




$p_3$




$p_4$




$p_5$




$p_6$



$p_7$



# The density of a five-step random walk



$$p_5(x) = \int_0^\infty xtJ_0(xt)J_0^5(t) dt$$

“... the graphical construction, however carefully reinvestigated, did not permit of our considering the curve to be anything but a **straight line**. . . Even if it is not absolutely true, it exemplifies the extraordinary power of such integrals of  $J$  products to give extremely close approximations to such simple forms as horizontal lines.”

**Karl Pearson, 1906**



**H. E. Fettis**

*On a conjecture of Karl Pearson*  
Rider Anniversary Volume, p. 39–54, 1963

$$p_2(x) = \frac{2}{\pi\sqrt{4-x^2}}$$

easy

$$p_3(x) = \operatorname{Re} \left( \frac{\sqrt{x}}{\pi^2} K \left( \sqrt{\frac{(x+1)^3(3-x)}{16x}} \right) \right)$$

G. J. Bennett  
1905

$$p_4(x) = ??$$

⋮

$$p_n(x) = \int_0^\infty xt J_0(xt) J_0^n(t) dt$$

J. C. Kluyver  
1906

# Classical results on the densities

$$p_2(x) = \frac{2}{\pi\sqrt{4-x^2}}$$

easy

$$p_3(x) = \operatorname{Re} \left( \frac{\sqrt{x}}{\pi^2} K \left( \sqrt{\frac{(x+1)^3(3-x)}{16x}} \right) \right)$$


G. J. Bennett  
1905

$$p_4(x) = ??$$




⋮

$$p_n(x) = \int_0^\infty xt J_0(xt) J_0^n(t) dt$$

J. C. Kluyver  
1906



# (New) Hypergeometric formulae



$$p_2(x) = \frac{2}{\pi\sqrt{4-x^2}}$$

easy

$$p_3(x) = \frac{2\sqrt{3}}{\pi} \frac{x}{(3+x^2)} {}_2F_1\left(\begin{matrix} \frac{1}{3}, \frac{2}{3} \\ 1 \end{matrix} \middle| \frac{x^2(9-x^2)^2}{(3+x^2)^3}\right)$$

classical  
with a spin

$$p_4(x) = \frac{2}{\pi^2} \frac{\sqrt{16-x^2}}{x} \operatorname{Re} {}_3F_2\left(\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ \frac{5}{6}, \frac{7}{6} \end{matrix} \middle| \frac{(16-x^2)^3}{108x^4}\right)$$

new  
BSWZ

- $s$ th moment  $W_n(s)$  of the density  $p_n$ :

$$W_n(s) = \int_0^\infty x^s p_n(x) dx = \int_{[0,1]^n} \left| \sum_{k=1}^n e^{2\pi x_k i} \right|^s d\mathbf{x}$$

# Moments of random walks

- $s$ th moment  $W_n(s)$  of the density  $p_n$ :

$$W_n(s) = \int_0^\infty x^s p_n(x) dx = \int_{[0,1]^n} \left| \sum_{k=1}^n e^{2\pi x_k i} \right|^s dx$$

THM  
Borwein-  
Nuyens-  
S-Wan  
2010

$$W_n(2k) = \sum_{a_1 + \dots + a_n = k} \binom{k}{a_1, \dots, a_n}^2$$

- $W_n(2k)$  counts the number of **abelian squares**: strings  $xy$  of length  $2k$  from an alphabet with  $n$  letters such that  $y$  is a permutation of  $x$ .
- Introduced by Erdős and studied by others.

EG

$$W_2(2k) = \binom{2k}{k}$$

# Even moments

$n$	$s = 0$	$s = 2$	$s = 4$	$s = 6$	$s = 8$	$s = 10$	Sloane's
2	1	2	6	20	70	252	A000984
3	1	3	15	93	639	4653	A002893
4	1	4	28	256	2716	31504	A002895
5	1	5	45	545	7885	127905	A169714
6	1	6	66	996	18306	384156	A169715

EG

$$W_3(2k) = \sum_{j=0}^k \binom{k}{j}^2 \binom{2j}{j}$$

Apéry-like

$$W_4(2k) = \sum_{j=0}^k \binom{k}{j}^2 \binom{2j}{j} \binom{2(k-j)}{k-j}$$

Domb numbers

# Even moments

$n$	$s = 0$	$s = 2$	$s = 4$	$s = 6$	$s = 8$	$s = 10$	Sloane's
2	1	2	6	20	70	252	A000984
3	1	3	15	93	639	4653	A002893
4	1	4	28	256	2716	31504	A002895
5	1	5	45	545	7885	127905	A169714
6	1	6	66	996	18306	384156	A169715

EG

$$W_3(2k) = \sum_{j=0}^k \binom{k}{j}^2 \binom{2j}{j}$$

Apéry-like

$$W_4(2k) = \sum_{j=0}^k \binom{k}{j}^2 \binom{2j}{j} \binom{2(k-j)}{k-j}$$

Domb numbers

- Inevitable **recursions**

$$K \cdot f(k) = f(k+1)$$



$$[(k+2)^2 K^2 - (10k^2 + 30k + 23)K + 9(k+1)^2] \cdot W_3(2k) = 0$$

$$[(k+2)^3 K^2 - (2k+3)(10k^2 + 30k + 24)K + 64(k+1)^3] \cdot W_4(2k) = 0$$

- Via **Carlson's Theorem** these become functional equations

# Complex moments



- Analytic continuations:



- $W_3(s)$  has a simple pole at  $-2$  with residue  $\frac{2}{\sqrt{3}\pi}$

# Complex moments

- Analytic continuations:



- $W_3(s)$  has a simple pole at  $-2$  with residue  $\frac{2}{\sqrt{3}\pi}$





$W_3(s)$  has simple poles at  $-2k - 2$  with residue

$$\frac{2}{\pi\sqrt{3}} \frac{W_3(2k)}{3^{2k}}$$

# Complex moments

- Analytic continuations:



- $W_3(s)$  has a simple pole at  $-2$  with residue  $\frac{2}{\sqrt{3}\pi}$



$W_3(s)$  has simple poles at  $-2k - 2$  with residue


$$\frac{2}{\pi\sqrt{3}} \frac{W_3(2k)}{3^{2k}}$$





$W_4(s)$  has double poles at  $-2k - 2$  with lowest-order term

$$\frac{3}{2\pi^2} \frac{W_4(2k)}{8^{2k}}$$

# $W_4(s)$ in the complex plane



# $W_4(s)$ in the complex plane



Experimental and  
computational  
mathematics:  
Selected writings

Jonathan Borwein  
and  
Peter Borwein

PSIpress

- Mellin transform  $F(s)$  of  $f(x)$ :

$$\mathcal{M}[f; s] = \int_0^{\infty} x^s f(x) \frac{dx}{x}$$

$$W_n(s-1) = \mathcal{M}[p_n; s]$$

- Mellin transform  $F(s)$  of  $f(x)$ :

$$\mathcal{M}[f; s] = \int_0^{\infty} x^s f(x) \frac{dx}{x}$$

- $F(s)$  is analytic in a strip
- Functional properties:
  - $\mathcal{M}[x^\mu f(x); s] = F(s + \mu)$
  - $\mathcal{M}[D_x f(x); s] = -(s - 1)F(s - 1)$
  - $\mathcal{M}[-\theta_x f(x); s] = sF(s)$

$$W_n(s - 1) = \mathcal{M}[p_n; s]$$

Thus functional equations for  $F(s)$  translate into DEs for  $f(x)$

- Mellin transform  $F(s)$  of  $f(x)$ :

$$\mathcal{M}[f; s] = \int_0^{\infty} x^s f(x) \frac{dx}{x}$$

- $F(s)$  is analytic in a strip
- Functional properties:
  - $\mathcal{M}[x^\mu f(x); s] = F(s + \mu)$
  - $\mathcal{M}[D_x f(x); s] = -(s - 1)F(s - 1)$
  - $\mathcal{M}[-\theta_x f(x); s] = sF(s)$

- Poles of  $F(s)$  left of strip  $\implies$  asymptotics of  $f(x)$  at zero


$$\frac{1}{(s+m)^{n+1}}$$

$$W_n(s - 1) = \mathcal{M}[p_n; s]$$

Thus functional equations for  $F(s)$  translate into DEs for  $f(x)$

$$\frac{(-1)^n}{n!} x^m (\log x)^n$$

# The differential equation for $p_4$




$$[(s+4)^3 S^4 - 4(s+3)(5s^2 + 30s + 48)S^2 + 64(s+2)^3] \cdot W_4(s) = 0$$

translates into  $A_4 \cdot p_4(x) = 0$  with

$$A_4 = x^4(\theta_x + 1)^3 - 4x^2\theta_x(5\theta_x^2 + 3) + 64(\theta_x - 1)^3$$

# The differential equation for $p_4$




$$[(s+4)^3 S^4 - 4(s+3)(5s^2 + 30s + 48)S^2 + 64(s+2)^3] \cdot W_4(s) = 0$$

translates into  $A_4 \cdot p_4(x) = 0$  with

$$\begin{aligned} A_4 &= x^4(\theta_x + 1)^3 - 4x^2\theta_x(5\theta_x^2 + 3) + 64(\theta_x - 1)^3 \\ &= (x-4)(x-2)x^3(x+2)(x+4)D_x^3 + 6x^4(x^2-10)D_x^2 \\ &\quad + x(7x^4 - 32x^2 + 64)D_x + (x^2-8)(x^2+8) \end{aligned}$$

# The differential equation for $p_4$



$$\left[ (s+4)^3 S^4 - 4(s+3)(5s^2 + 30s + 48)S^2 + 64(s+2)^3 \right] \cdot W_4(s) = 0$$

translates into  $A_4 \cdot p_4(x) = 0$  with

$$\begin{aligned} A_4 &= x^4(\theta_x + 1)^3 - 4x^2\theta_x(5\theta_x^2 + 3) + 64(\theta_x - 1)^3 \\ &= (x-4)(x-2)x^3(x+2)(x+4)D_x^3 + 6x^4(x^2-10)D_x^2 \\ &\quad + x(7x^4 - 32x^2 + 64)D_x + (x^2-8)(x^2+8) \end{aligned}$$

**THM**  
Borwein-  
S-Wan-  
Zudilin,  
2011

- The density  $p_n$  satisfies a DE of order  $n-1$ .
- $p_n$  is real analytic except at 0 and the integers  $n, n-2, n-4, \dots$

$$s_{4,k} = \frac{3}{2\pi^2} \frac{W_4(2k)}{8^{2k}}$$

- $W_4(s)$  has double poles:

$$W_4(s) = \frac{s_{4,k}}{(s + 2k + 2)^2} + \frac{r_{4,k}}{s + 2k + 2} + O(1) \quad \text{as } s \rightarrow -2k - 2$$

$$p_4(x) = \sum_{k=0}^{\infty} (r_{4,k} - s_{4,k} \log(x)) x^{2k+1} \quad \text{for small } x \geq 0$$

$$s_{4,k} = \frac{3}{2\pi^2} \frac{W_4(2k)}{8^{2k}}$$

- $W_4(s)$  has double poles:

$$W_4(s) = \frac{s_{4,k}}{(s + 2k + 2)^2} + \frac{r_{4,k}}{s + 2k + 2} + O(1) \quad \text{as } s \rightarrow -2k - 2$$

$$p_4(x) = \sum_{k=0}^{\infty} (r_{4,k} - s_{4,k} \log(x)) x^{2k+1} \quad \text{for small } x \geq 0$$

- $y_0(z) := \sum_{k \geq 0} W_4(2k) z^k$  is the analytic solution of

$$[64z^2(\theta + 1)^3 - 2z(2\theta + 1)(5\theta^2 + 5\theta + 2) + \theta^3] \cdot y(z) = 0. \quad (\text{DE})$$

$$s_{4,k} = \frac{3}{2\pi^2} \frac{W_4(2k)}{8^{2k}}$$

- $W_4(s)$  has double poles:


$$W_4(s) = \frac{s_{4,k}}{(s + 2k + 2)^2} + \frac{r_{4,k}}{s + 2k + 2} + O(1) \quad \text{as } s \rightarrow -2k - 2$$

$$p_4(x) = \sum_{k=0}^{\infty} (r_{4,k} - s_{4,k} \log(x)) x^{2k+1} \quad \text{for small } x \geq 0$$

- $y_0(z) := \sum_{k \geq 0} W_4(2k) z^k$  is the analytic solution of

$$[64z^2(\theta + 1)^3 - 2z(2\theta + 1)(5\theta^2 + 5\theta + 2) + \theta^3] \cdot y(z) = 0. \quad (\text{DE})$$


- Let  $y_1(z)$  solve (DE) and  $y_1(z) - y_0(z) \log(z) \in z\mathbb{Q}[[z]]$ .  
Then  $p_4(x) = -\frac{3x}{4\pi^2} y_1(x^2/64)$ .



**THM**  
Chan-  
Chan-Liu  
2004;  
Rogers  
2009

Generating function for Domb numbers:

$$\sum_{k=0}^{\infty} W_4(2k)z^k = \frac{1}{1-4z} {}_3F_2 \left( \begin{matrix} \frac{1}{3}, \frac{1}{2}, \frac{2}{3} \\ 1, 1 \end{matrix} \middle| \frac{108z^2}{(1-4z)^3} \right)$$




**THM**  
Chan-  
Chan-Liu  
2004;  
Rogers  
2009

Generating function for Domb numbers:

$$\sum_{k=0}^{\infty} W_4(2k)z^k = \frac{1}{1-4z} {}_3F_2 \left( \begin{matrix} \frac{1}{3}, \frac{1}{2}, \frac{2}{3} \\ 1, 1 \end{matrix} \middle| \frac{108z^2}{(1-4z)^3} \right)$$

- Basis at  $\infty$  for the hypergeometric equation of  ${}_3F_2 \left( \frac{1}{3}, \frac{1}{2}, \frac{2}{3} \middle| t \right)$ :  
[as  $x \rightarrow 4$  then  $z = \frac{x^2}{64} \rightarrow \frac{1}{4}$  and  $t = \frac{108z^2}{(1-4z)^3} \rightarrow \infty$ ]

$$t^{-1/3} {}_3F_2 \left( \begin{matrix} \frac{1}{3}, \frac{1}{3}, \frac{1}{3} \\ \frac{2}{3}, \frac{5}{6} \end{matrix} \middle| \frac{1}{t} \right), \quad t^{-1/2} {}_3F_2 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ \frac{5}{6}, \frac{7}{6} \end{matrix} \middle| \frac{1}{t} \right), \quad t^{-2/3} {}_3F_2 \left( \begin{matrix} \frac{2}{3}, \frac{2}{3}, \frac{2}{3} \\ \frac{4}{3}, \frac{7}{6} \end{matrix} \middle| \frac{1}{t} \right)$$



**THM** Generating function for Domb numbers:

Chan-  
Chan-Liu  
2004;  
Rogers  
2009

$$\sum_{k=0}^{\infty} W_4(2k)z^k = \frac{1}{1-4z} {}_3F_2 \left( \begin{matrix} \frac{1}{3}, \frac{1}{2}, \frac{2}{3} \\ 1, 1 \end{matrix} \middle| \frac{108z^2}{(1-4z)^3} \right)$$


- Basis at  $\infty$  for the hypergeometric equation of  ${}_3F_2 \left( \frac{1}{3}, \frac{1}{2}, \frac{2}{3} \middle| t \right)$ :  
[as  $x \rightarrow 4$  then  $z = \frac{x^2}{64} \rightarrow \frac{1}{4}$  and  $t = \frac{108z^2}{(1-4z)^3} \rightarrow \infty$ ]

$$t^{-1/3} {}_3F_2 \left( \begin{matrix} \frac{1}{3}, \frac{1}{3}, \frac{1}{3} \\ \frac{2}{3}, \frac{5}{6} \end{matrix} \middle| \frac{1}{t} \right), \quad t^{-1/2} {}_3F_2 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ \frac{5}{6}, \frac{7}{6} \end{matrix} \middle| \frac{1}{t} \right), \quad t^{-2/3} {}_3F_2 \left( \begin{matrix} \frac{2}{3}, \frac{2}{3}, \frac{2}{3} \\ \frac{4}{3}, \frac{7}{6} \end{matrix} \middle| \frac{1}{t} \right)$$

**THM** For  $2 \leq x \leq 4$ ,

Borwein-  
S-Wan-  
Zudilin  
2011

$$p_4(x) = \frac{2}{\pi^2} \frac{\sqrt{16-x^2}}{x} {}_3F_2 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ \frac{5}{6}, \frac{7}{6} \end{matrix} \middle| \frac{(16-x^2)^3}{108x^4} \right).$$



**THM** Chan-Chan-Liu 2004; Rogers 2009

Generating function for Domb numbers:

$$\sum_{k=0}^{\infty} W_4(2k)z^k = \frac{1}{1-4z} {}_3F_2 \left( \begin{matrix} \frac{1}{3}, \frac{1}{2}, \frac{2}{3} \\ 1, 1 \end{matrix} \middle| \frac{108z^2}{(1-4z)^3} \right)$$

- Basis at  $\infty$  for the hypergeometric equation of  ${}_3F_2 \left( \frac{1}{3}, \frac{1}{2}, \frac{2}{3} \middle| t \right)$ :  
[as  $x \rightarrow 4$  then  $z = \frac{x^2}{64} \rightarrow \frac{1}{4}$  and  $t = \frac{108z^2}{(1-4z)^3} \rightarrow \infty$ ]

$$t^{-1/3} {}_3F_2 \left( \begin{matrix} \frac{1}{3}, \frac{1}{3}, \frac{1}{3} \\ \frac{2}{3}, \frac{5}{6} \end{matrix} \middle| \frac{1}{t} \right), \quad t^{-1/2} {}_3F_2 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ \frac{5}{6}, \frac{7}{6} \end{matrix} \middle| \frac{1}{t} \right), \quad t^{-2/3} {}_3F_2 \left( \begin{matrix} \frac{2}{3}, \frac{2}{3}, \frac{2}{3} \\ \frac{4}{3}, \frac{7}{6} \end{matrix} \middle| \frac{1}{t} \right)$$


**THM** Borwein-S-Wan-Zudilin 2011

For  ~~$2 \leq x \leq 4$~~   $0 \leq x \leq 4$ ,

$$p_4(x) = \mathbf{Re} \frac{2}{\pi^2} \frac{\sqrt{16-x^2}}{x} {}_3F_2 \left( \begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ \frac{5}{6}, \frac{7}{6} \end{matrix} \middle| \frac{(16-x^2)^3}{108x^4} \right).$$

## The density of a five-step random walk, again

$$p_5(x) = 0.32993x + 0.0066167x^3 + 0.00026233x^5 + 0.000014119x^7 + O(x^9)$$




$$p_5(x) = \int_0^\infty xtJ_0(xt)J_0^5(t) dt$$

“ ... the graphical construction, however carefully reinvestigated, did not permit of our considering the curve to be anything but a **straight line**. . . Even if it is not absolutely true, it exemplifies the extraordinary power of such integrals of  $J$  products to give extremely close approximations to such simple forms as horizontal lines.

Karl Pearson, 1906

## The density of a five-step random walk, again

$$p_5(x) = \underbrace{0.32993}_{=p_4(1)}x + 0.0066167x^3 + 0.00026233x^5 + 0.000014119x^7 + O(x^9)$$



$$p_5(x) = \int_0^\infty xtJ_0(xt)J_0^5(t) dt$$

“... the graphical construction, however carefully reinvestigated, did not permit of our considering the curve to be anything but a **straight line**. . . Even if it is not absolutely true, it exemplifies the extraordinary power of such integrals of  $J$  products to give extremely close approximations to such simple forms as horizontal lines.”

Karl Pearson, 1906

- THM** Let  $f(\tau)$  be a modular form and  $x(\tau)$  a modular function w.r.t.  $\Gamma$ .
- Then  $y(x)$  defined by  $f(\tau) = y(x(\tau))$  satisfies a linear DE.
  - If  $x(\tau)$  is a Hauptmodul for  $\Gamma$ , then the DE has polynomial coefficients.
  - The solutions of the DE are  $y(x), \tau y(x), \tau^2 y(x), \dots$

**THM** Let  $f(\tau)$  be a modular form and  $x(\tau)$  a modular function w.r.t.  $\Gamma$ .

- Then  $y(x)$  defined by  $f(\tau) = y(x(\tau))$  satisfies a linear DE.
- If  $x(\tau)$  is a Hauptmodul for  $\Gamma$ , then the DE has polynomial coefficients.
- The solutions of the DE are  $y(x), \tau y(x), \tau^2 y(x), \dots$

**EG**  
Classic

$${}_2F_1 \left( \begin{matrix} 1/2, 1/2 \\ 1 \end{matrix} \middle| \lambda(\tau) \right) = \theta_3(\tau)^2$$

- $\lambda(\tau) = 16 \frac{\eta(\tau/2)^8 \eta(2\tau)^{16}}{\eta(\tau)^{24}}$  is the elliptic lambda function, a Hauptmodul for  $\Gamma(2)$ .
- $\theta_3(\tau) = \frac{\eta(\tau)^5}{\eta(\tau/2)^2 \eta(2\tau)^2}$  is the usual Jacobi theta function.

**THM** Let  $f(\tau)$  be a modular form and  $x(\tau)$  a modular function w.r.t.  $\Gamma$ .

- Then  $y(x)$  defined by  $f(\tau) = y(x(\tau))$  satisfies a linear DE.
- If  $x(\tau)$  is a Hauptmodul for  $\Gamma$ , then the DE has polynomial coefficients.
- The solutions of the DE are  $y(x), \tau y(x), \tau^2 y(x), \dots$

**EG**  
Chan-  
Chan-Liu  
2004

$$\begin{aligned}x(\tau) &= - \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^6, & f(\tau) &= \frac{(\eta(\tau)\eta(3\tau))^4}{(\eta(2\tau)\eta(6\tau))^2} \\ &= -q - 6q^2 - 21q^3 - 68q^4 + \dots & &= 1 - 4q + 4q^2 - 4q^3 + 20q^4 + \dots\end{aligned}$$

Here,  $\Gamma = \left\langle \Gamma_0(6), \frac{1}{\sqrt{3}} \begin{pmatrix} 3 & -2 \\ 6 & -3 \end{pmatrix} \right\rangle$ .

# Modular differential equations

**THM** Let  $f(\tau)$  be a modular form and  $x(\tau)$  a modular function w.r.t.  $\Gamma$ .

- Then  $y(x)$  defined by  $f(\tau) = y(x(\tau))$  satisfies a linear DE.
- If  $x(\tau)$  is a Hauptmodul for  $\Gamma$ , then the DE has polynomial coefficients.
- The solutions of the DE are  $y(x), \tau y(x), \tau^2 y(x), \dots$

**EG**  
Chan-  
Liu  
2004

$$\begin{aligned} x(\tau) &= - \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^6, & f(\tau) &= \frac{(\eta(\tau)\eta(3\tau))^4}{(\eta(2\tau)\eta(6\tau))^2} \\ &= -q - 6q^2 - 21q^3 - 68q^4 + \dots & &= 1 - 4q + 4q^2 - 4q^3 + 20q^4 + \dots \end{aligned}$$

Here,  $\Gamma = \left\langle \Gamma_0(6), \frac{1}{\sqrt{3}} \begin{pmatrix} 3 & -2 \\ 6 & -3 \end{pmatrix} \right\rangle$ . Then, in a neighborhood of  $i\infty$ ,

$$f(\tau) = y_0(x(\tau)) = \sum_{k \geq 0} W_4(2k) x(\tau)^k.$$

For  $\tau = -1/2 + iy$  and  $y > 0$ :

$$p_4 \left( \underbrace{8i \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^3}_{=\sqrt{64x(\tau)}} \right) = \frac{6(2\tau + 1)}{\pi} \underbrace{\eta(\tau)\eta(2\tau)\eta(3\tau)\eta(6\tau)}_{=\sqrt{-x(\tau)}f(\tau)}$$

**THM**  
Borwein-  
S-Wan-  
Zudilin  
2011

For  $\tau = -1/2 + iy$  and  $y > 0$ :

$$p_4 \left( \underbrace{8i \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^3}_{=\sqrt{64x(\tau)}} \right) = \frac{6(2\tau + 1)}{\pi} \underbrace{\eta(\tau)\eta(2\tau)\eta(3\tau)\eta(6\tau)}_{=\sqrt{-x(\tau)}f(\tau)}$$

- When  $\tau = -\frac{1}{2} + \frac{1}{6}\sqrt{-15}$ , one obtains  $p_4(1) = p'_5(0)$  as an  $\eta$ -product.

# Modular parametrization of $p_4$

**THM**  
Borwein-  
S-Wan-  
Zudilin  
2011

For  $\tau = -1/2 + iy$  and  $y > 0$ :

$$p_4 \left( \underbrace{8i \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^3}_{=\sqrt{64x(\tau)}} \right) = \frac{6(2\tau + 1)}{\pi} \underbrace{\eta(\tau)\eta(2\tau)\eta(3\tau)\eta(6\tau)}_{=\sqrt{-x(\tau)}f(\tau)}$$

- When  $\tau = -\frac{1}{2} + \frac{1}{6}\sqrt{-15}$ , one obtains  $p_4(1) = p'_5(0)$  as an  $\eta$ -product.
- Applying the Chowla–Selberg formula, eventually leads to:

**COR**

$$p_4(1) = p'_5(0) = \frac{\sqrt{5}}{40\pi^4} \Gamma\left(\frac{1}{15}\right)\Gamma\left(\frac{2}{15}\right)\Gamma\left(\frac{4}{15}\right)\Gamma\left(\frac{8}{15}\right) \approx 0.32993$$

# Modular parametrization of $p_4$

**THM**  
Borwein-  
S-Wan-  
Zudilin  
2011

For  $\tau = -1/2 + iy$  and  $y > 0$ :

$$p_4 \left( \underbrace{8i \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^3}_{=\sqrt{64x(\tau)}} \right) = \frac{6(2\tau + 1)}{\pi} \underbrace{\eta(\tau)\eta(2\tau)\eta(3\tau)\eta(6\tau)}_{=\sqrt{-x(\tau)}f(\tau)}$$

- When  $\tau = -\frac{1}{2} + \frac{1}{6}\sqrt{-15}$ , one obtains  $p_4(1) = p'_5(0)$  as an  $\eta$ -product.
- Applying the Chowla–Selberg formula, eventually leads to:

**COR**

$$p_4(1) = p'_5(0) = \frac{\sqrt{5}}{40\pi^4} \Gamma\left(\frac{1}{15}\right)\Gamma\left(\frac{2}{15}\right)\Gamma\left(\frac{4}{15}\right)\Gamma\left(\frac{8}{15}\right) \approx 0.32993$$

**Fact**

If  $\sigma_1, \sigma_2 \in \mathcal{H}$  both belong to  $\mathbb{Q}(\sqrt{-d})$ , then the quotient  $\eta(\sigma_1) / \eta(\sigma_2)$  is an algebraic number.

# Chowla–Selberg formula

THM  
Chowla–  
Selberg  
1967

$$\prod_{j=1}^h a_j^{-6} |\eta(\tau_j)|^{24} = \frac{1}{(2\pi|d|)^{6h}} \left[ \prod_{k=1}^{|d|} \Gamma\left(\frac{k}{|d|}\right)^{\binom{|d|}{k}} \right]^{3w}$$

where the product is over reduced binary quadratic forms  $[a_j, b_j, c_j]$  of discriminant  $d < 0$ .

$$\tau_j = \frac{-b_j + \sqrt{d}}{2a_j}$$

# Chowla–Selberg formula

**THM**  
Chowla–  
Selberg  
1967

$$\prod_{j=1}^h a_j^{-6} |\eta(\tau_j)|^{24} = \frac{1}{(2\pi|d|)^{6h}} \left[ \prod_{k=1}^{|d|} \Gamma\left(\frac{k}{|d|}\right)^{\binom{d}{k}} \right]^{3w}$$

where the product is over reduced binary quadratic forms  $[a_j, b_j, c_j]$  of discriminant  $d < 0$ .


$$\tau_j = \frac{-b_j + \sqrt{d}}{2a_j}$$

**EG**  $\mathbb{Q}(\sqrt{-15})$  has discriminant  $\Delta = -15$  and class number  $h = 2$ .

$$\begin{aligned} Q_1 &= [1, 1, 4] & Q_2 &= [2, 1, 2] \\ \tau_1 &= -\frac{1}{2} + \frac{1}{2}\sqrt{-15}, & \tau_2 &= \frac{1}{2}\tau_1 \end{aligned}$$

$$\begin{aligned} \frac{1}{\sqrt{2}} |\eta(\tau_1)\eta(\tau_2)|^2 &= \frac{1}{30\pi} \left( \frac{\Gamma(\frac{1}{15})\Gamma(\frac{2}{15})\Gamma(\frac{4}{15})\Gamma(\frac{8}{15})}{\Gamma(\frac{7}{15})\Gamma(\frac{11}{15})\Gamma(\frac{13}{15})\Gamma(\frac{14}{15})} \right)^{1/2} \\ &= \frac{1}{120\pi^3} \Gamma\left(\frac{1}{15}\right)\Gamma\left(\frac{2}{15}\right)\Gamma\left(\frac{4}{15}\right)\Gamma\left(\frac{8}{15}\right) \end{aligned}$$

# What we know about $p_5$



- $W_5(s)$  has simple poles at  $-2k - 2$  with residue  $r_{5,k}$
- Hence:  $p_5(x) = \sum_{k=0}^{\infty} r_{5,k} x^{2k+1}$

**THM**  
Borwein-  
S-Wan-  
Zudilin,  
2011

Surprising bonus of the modularity of  $p_4$ :

$$r_{5,0} = p_4(1) = \frac{\sqrt{5}}{40} \frac{\Gamma(\frac{1}{15})\Gamma(\frac{2}{15})\Gamma(\frac{4}{15})\Gamma(\frac{8}{15})}{\pi^4}$$
$$r_{5,1} \stackrel{?}{=} \frac{13}{225} r_{5,0} - \frac{2}{5\pi^4} \frac{1}{r_{5,0}}$$

- Other residues given recursively
- $p_5$  solves the DE

$$\left[ x^6(\theta + 1)^4 - x^4(35\theta^4 + 42\theta^2 + 3) + x^2(259(\theta - 1)^4 + 104(\theta - 1)^2) - (15(\theta - 3)(\theta - 1))^2 \right] \cdot p_5(x) = 0$$

**DEF** (Logarithmic) Mahler measure of  $p(x_1, \dots, x_n)$ :

$$\mu(p) := \int_0^1 \cdots \int_0^1 \log |p(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| dt_1 dt_2 \dots dt_n$$

**DEF** (Logarithmic) Mahler measure of  $p(x_1, \dots, x_n)$ :

$$\mu(p) := \int_0^1 \cdots \int_0^1 \log |p(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| dt_1 dt_2 \dots dt_n$$

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$

**EG**

$$W_n'(0) = \mu(x_1 + \dots + x_n) = \mu(1 + x_1 + \dots + x_{n-1})$$

**DEF** (Logarithmic) Mahler measure of  $p(x_1, \dots, x_n)$ :

$$\mu(p) := \int_0^1 \cdots \int_0^1 \log |p(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| dt_1 dt_2 \dots dt_n$$

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$

**EG**

$$W_n'(0) = \mu(x_1 + \dots + x_n) = \mu(1 + x_1 + \dots + x_{n-1})$$

**EG**  
Smyth,  
1981

$$\begin{aligned} \mu(1 + x + y) &= \frac{3\sqrt{3}}{4\pi} L(\chi_{-3}, 2) &&= W_3'(0) \\ \mu(1 + x + y + z) &= \frac{7}{2} \frac{\zeta(3)}{\pi^2} &&= W_4'(0) \end{aligned}$$

$$L(\chi_{-3}, s) = 1 - \frac{1}{2^s} + \frac{1}{4^s} - \frac{1}{5^s} + \frac{1}{7^s} - \dots$$

**EG**  
Rogers–  
Zudilin,  
2011

Typical conjecture (Deninger, 1997):

$$\mu(1 + x + y + 1/x + 1/y) = \left( \frac{\sqrt{-15}}{2\pi i} \right)^2 L(f_{15}, 2) = L'(f_{15}, 0)$$

where  $f_{15}$  is associated with an elliptic curve of conductor 15.

**EG**  
Rogers-  
Zudilin,  
2011

Typical conjecture (Deninger, 1997):

$$\mu(1 + x + y + 1/x + 1/y) = \left( \frac{\sqrt{-15}}{2\pi i} \right)^2 L(f_{15}, 2) = L'(f_{15}, 0)$$

where  $f_{15}$  is associated with an elliptic curve of conductor 15.

**CONJ**  
Rodriguez-  
Villegas

$$W'_5(0) \stackrel{?}{=} \left( \frac{\sqrt{-15}}{2\pi i} \right)^5 3! L(g_{15}, 4) = -L'(g_{15}, -1)$$

where  $g_{15} = \eta(3\tau)^3 \eta(5\tau)^3 + \eta(\tau)^3 \eta(15\tau)^3$  (weight 3, level 15).

**EG**  
Rogers–  
Zudilin,  
2011

Typical conjecture (Deninger, 1997):

$$\mu(1 + x + y + 1/x + 1/y) = \left( \frac{\sqrt{-15}}{2\pi i} \right)^2 L(f_{15}, 2) = L'(f_{15}, 0)$$

where  $f_{15}$  is associated with an elliptic curve of conductor 15.

**CONJ**  
Rodríguez-  
Villegas

$$W'_5(0) \stackrel{?}{=} \left( \frac{\sqrt{-15}}{2\pi i} \right)^5 3! L(g_{15}, 4) = -L'(g_{15}, -1)$$

where  $g_{15} = \eta(3\tau)^3 \eta(5\tau)^3 + \eta(\tau)^3 \eta(15\tau)^3$  (weight 3, level 15).

**CONJ**  
Rodríguez-  
Villegas

$$W'_6(0) \stackrel{?}{=} 8 \left( \frac{\sqrt{-6}}{2\pi i} \right)^6 4! L(g_6, 5) = -8L'(g_6, -1)$$

where  $g_6 = \eta(\tau)^2 \eta(2\tau)^2 \eta(3\tau)^2 \eta(6\tau)^2$  (weight 4, level 6).

EG  
Chan-  
Chan-Liu  
2004

$$\begin{aligned}x(\tau) &= - \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^6, & f(\tau) &= \frac{(\eta(\tau)\eta(3\tau))^4}{(\eta(2\tau)\eta(6\tau))^2} \\ &= -q - 6q^2 - 21q^3 - 68q^4 + \dots & &= 1 - 4q + 4q^2 - 4q^3 + 20q^4 + \dots\end{aligned}$$

$$f(\tau) = y_0(x(\tau)) = \sum_{k \geq 0} W_4(2k)x(\tau)^k.$$

EG  
Chan-  
Chan-Liu  
2004

$$\begin{aligned}x(\tau) &= - \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^6, & f(\tau) &= \frac{(\eta(\tau)\eta(3\tau))^4}{(\eta(2\tau)\eta(6\tau))^2} \\ &= -q - 6q^2 - 21q^3 - 68q^4 + \dots & &= 1 - 4q + 4q^2 - 4q^3 + 20q^4 + \dots\end{aligned}$$

$$f(\tau) = y_0(x(\tau)) = \sum_{k \geq 0} W_4(2k)x(\tau)^k.$$

- Double  $L$ -function:

$$f = \sum a_n q^n, \quad g = \sum b_n q^n$$

$$L(f, g, s, t) \stackrel{\text{“}=\text{”}}{=} \sum_{n \geq 1} \sum_{m \geq 0} \frac{a_n b_m}{n^s (n+m)^t}$$

EG  
Chan-  
Liu  
2004

$$\begin{aligned}x(\tau) &= - \left( \frac{\eta(2\tau)\eta(6\tau)}{\eta(\tau)\eta(3\tau)} \right)^6, & f(\tau) &= \frac{(\eta(\tau)\eta(3\tau))^4}{(\eta(2\tau)\eta(6\tau))^2} \\ &= -q - 6q^2 - 21q^3 - 68q^4 + \dots & &= 1 - 4q + 4q^2 - 4q^3 + 20q^4 + \dots\end{aligned}$$

$$f(\tau) = y_0(x(\tau)) = \sum_{k \geq 0} W_4(2k)x(\tau)^k.$$

- Double  $L$ -function:

$$f = \sum a_n q^n, \quad g = \sum b_n q^n$$

$$L(f, g, s, t) \stackrel{“=”}{=} \sum_{n \geq 1} \sum_{m \geq 0} \frac{a_n b_m}{n^s (n+m)^t}$$

THM  
Shinder-  
Vlasenko  
2012

$$W'_5(0) - \frac{4}{5}W'_4(0) = \frac{3\sqrt{5}\Omega_{15}^2}{20\pi} L(g_3, g_1, 3, 1) - \frac{3\sqrt{5}}{10\pi^3\Omega_{15}^2} L(g_2, g_1, 3, 1)$$

where

$$g_1 = \frac{Dx}{x} f, \quad g_2 = \frac{x}{1-x} g_1, \quad g_3 = \frac{x(212x^2 + 251x - 13)}{(1-x)^3} g_1.$$

## The Shinder-Vlasenko starting point

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$
- $W'_n(0) = \frac{1}{2}\mu(p_n)$  where  $p_n = (1 + x_1 + \dots + x_{n-1})(1 + \frac{1}{x_1} + \dots + \frac{1}{x_{n-1}})$

# The Shinder-Vlasenko starting point

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$
- $W'_n(0) = \frac{1}{2}\mu(p_n)$  where  $p_n = (1 + x_1 + \dots + x_{n-1})(1 + \frac{1}{x_1} + \dots + \frac{1}{x_{n-1}})$

**Trick**  
Rodriguez-  
Villegas

$$\int_{[0,1]^n} \log \left[ p_n (e^{2\pi i t_1}, \dots, e^{2\pi i t_n}) - \frac{1}{\lambda} \right] dt$$

# The Shinder-Vlasenko starting point

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$
- $W'_n(0) = \frac{1}{2}\mu(p_n)$  where  $p_n = (1 + x_1 + \dots + x_{n-1})(1 + \frac{1}{x_1} + \dots + \frac{1}{x_{n-1}})$

**Trick**  
Rodriguez-  
Villegas

$$\begin{aligned} & \int_{[0,1]^n} \log \left[ p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n}) - \frac{1}{\lambda} \right] dt \\ &= -\log(-\lambda) - \sum_{k \geq 1} \frac{\lambda^k}{k} \int_{[0,1]^n} p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})^k dt \end{aligned}$$

# The Shinder-Vlasenko starting point

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$
- $W'_n(0) = \frac{1}{2}\mu(p_n)$  where  $p_n = (1 + x_1 + \dots + x_{n-1})(1 + \frac{1}{x_1} + \dots + \frac{1}{x_{n-1}})$

**Trick**  
Rodríguez-  
Villegas

$$\begin{aligned} & \int_{[0,1]^n} \log \left[ p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n}) - \frac{1}{\lambda} \right] dt \\ &= -\log(-\lambda) - \sum_{k \geq 1} \frac{\lambda^k}{k} \int_{[0,1]^n} p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})^k dt \\ &= -\log(-\lambda) - \sum_{k \geq 1} \frac{\lambda^k}{k} W_n(2k) \end{aligned}$$

# The Shinder-Vlasenko starting point

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$
- $W'_n(0) = \frac{1}{2}\mu(p_n)$  where  $p_n = (1 + x_1 + \dots + x_{n-1})(1 + \frac{1}{x_1} + \dots + \frac{1}{x_{n-1}})$

**Trick**  
Rodríguez-  
Villegas

$$\begin{aligned} & \int_{[0,1]^n} \log \left[ p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n}) - \frac{1}{\lambda} \right] dt \\ &= -\log(-\lambda) - \sum_{k \geq 1} \frac{\lambda^k}{k} \int_{[0,1]^n} p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})^k dt \\ &= -\log(-\lambda) - \sum_{k \geq 1} \frac{\lambda^k}{k} W_n(2k) = - \left[ \lambda \frac{d}{d\lambda} \right]^{-1} \sum_{k \geq 0} W_n(2k) \lambda^k \end{aligned}$$

# The Shinder-Vlasenko starting point

- $W_n(s) = \int_{[0,1]^n} |e^{2\pi x_1 i} + \dots + e^{2\pi x_n i}|^s d\mathbf{x}$
- $W'_n(0) = \frac{1}{2}\mu(p_n)$  where  $p_n = (1 + x_1 + \dots + x_{n-1})(1 + \frac{1}{x_1} + \dots + \frac{1}{x_{n-1}})$

**Trick**  
Rodríguez-  
Villegas

$$\begin{aligned} & \int_{[0,1]^n} \log \left[ p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n}) - \frac{1}{\lambda} \right] dt \\ &= -\log(-\lambda) - \sum_{k \geq 1} \frac{\lambda^k}{k} \int_{[0,1]^n} p_n(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})^k dt \\ &= -\log(-\lambda) - \sum_{k \geq 1} \frac{\lambda^k}{k} W_n(2k) = - \left[ \lambda \frac{d}{d\lambda} \right]^{-1} \sum_{k \geq 0} W_n(2k) \lambda^k \end{aligned}$$

Hence, analytically continuing along the negative real axis,

$$\mu(p_n) = -\operatorname{Re} \left[ \lambda \frac{d}{d\lambda} \right]^{-1} \sum_{k \geq 0} W_n(2k) \lambda^k \Big|_{\lambda=\infty}.$$

- The differential equations for  $n \geq 5$  are not modular.  
Can one profitably bring vector-valued modular forms into the picture?
- Given a linear differential equation **automatically** find its “hypergeometric-type” solutions.  
Promising work by Mark van Hoeij and his group
- More about the five step case? Average distance travelled?  
$$W_n(1) = n \int_0^\infty J_1(x) J_0(x)^{n-1} \frac{dx}{x}$$
- Countless generalizations ...  
higher dimensions, different step sizes, ...

# THANK YOU!

- Slides for this talk will be available from my website:  
<http://arminstraub.com/talks>



**J. Borwein, D. Nuyens, A. Straub, J. Wan**

*Some arithmetic properties of short random walk integrals*

The Ramanujan Journal, Vol. 26, Nr. 1, 2011, p. 109-132



**J. Borwein, A. Straub, J. Wan**

*Three-step and four-step random walk integrals*

Experimental Mathematics — to appear



**J. Borwein, A. Straub, J. Wan, W. Zudilin (appendix by D. Zagier)**

*Densities of short uniform random walks*

Canadian Journal of Mathematics — to appear

...

- Representations for  $W_n(s)$  give us, for instance,

$$\begin{aligned}W'_n(0) &= \log(2) - \gamma - \int_0^1 (J_0^n(x) - 1) \frac{dx}{x} - \int_1^\infty J_0^n(x) \frac{dx}{x} \\ &= \log(2) - \gamma - n \int_0^\infty \log(x) J_0^{n-1}(x) J_1(x) dx.\end{aligned}$$

**DEF** Multiple Mahler measure of polynomials  $p_i(x_1, \dots, x_n)$ :

$$\mu(p_1, \dots, p_k) := \int_{[0,1]^n} \prod_{i=1}^k \log |p_i(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| \, d\mathbf{t}$$

$$\mu_k(p) := \int_{[0,1]^n} \log^k |p(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| \, d\mathbf{t}$$

# (Multiple) Mahler measure

**DEF** Multiple Mahler measure of polynomials  $p_i(x_1, \dots, x_n)$ :

$$\mu(p_1, \dots, p_k) := \int_{[0,1]^n} \prod_{i=1}^k \log |p_i(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| dt$$

$$\mu_k(p) := \int_{[0,1]^n} \log^k |p(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| dt$$

**EG**

$$W_n^{(k)}(0) = \mu_k(1 + x_1 + \dots + x_{n-1})$$

# Moments of a 3-step random walk

EG

Borwein-  
Borwein-  
S-Wan

$$\mu_1(1+x+y) = \frac{3}{2\pi} \text{LS}_2 \left( \frac{2\pi}{3} \right)$$

$$\mu_2(1+x+y) = \frac{3}{\pi} \text{LS}_3 \left( \frac{2\pi}{3} \right) + \frac{\pi^2}{4}$$

$$\mu_3(1+x+y) \stackrel{?}{=} \frac{6}{\pi} \text{LS}_4 \left( \frac{2\pi}{3} \right) - \frac{9}{\pi} \text{Cl}_4 \left( \frac{\pi}{3} \right) \\ - \frac{\pi}{4} \text{Cl}_2 \left( \frac{\pi}{3} \right) - \frac{13}{2} \zeta(3)$$

$$\mu_4(1+x+y) \stackrel{?}{=} \frac{12}{\pi} \text{LS}_5 \left( \frac{2\pi}{3} \right) - \frac{49}{3\pi} \text{LS}_5 \left( \frac{\pi}{3} \right) + \frac{81}{\pi} \text{Gl}_{4,1} \left( \frac{2\pi}{3} \right) \\ + 3\pi \text{Gl}_{2,1} \left( \frac{2\pi}{3} \right) + \frac{2}{\pi} \zeta(3) \text{Cl}_2 \left( \frac{\pi}{3} \right) \\ + \text{Cl}_2 \left( \frac{\pi}{3} \right)^2 - \frac{29}{90} \pi^4$$

- Using the residues  $r_{5,k} = \text{Res}_{-2k-2} W_5$ :

$$p_5(x) = \sum_{k=0}^{\infty} r_{5,k} x^{2k+1}$$

EG

$$r_{5,0} = \frac{16 + 1140W_5'(0) - 804W_5'(2) + 64W_5'(4)}{225},$$
$$r_{5,1} = \frac{26r_{5,0} - 16 - 20W_5'(0) + 4W_5'(2)}{225}.$$

- Unfortunately, the Mahler measure  $W_5'(0)$  “cancels” out.