Midterm #3

MATH 126-102 — Calculus II
Thursday, Nov 19, 2026

Please print your name:

No notes, graphing calculators or other tools are permitted. There are 30 points in total. You need to show work to receive full credit.

Good luck!

Problem 1. (6 points) Compute the following series (or state that it diverges):

(a) $\displaystyle\sum_{n=1}^{\infty} \frac{5-3^n}{4^n}$

(b) $\displaystyle\sum_{n=1}^{\infty} \frac{5^n-3}{4^n}$

Solution.

(a) $\displaystyle\sum_{n=1}^{\infty} \frac{5-3^n}{4^n} = 5\sum_{n=1}^{\infty}\left(\frac{1}{4}\right)^n - \sum_{n=1}^{\infty}\left(\frac{3}{4}\right)^n = 5\left(\frac{1}{1-\frac{1}{4}} - \left(\frac{1}{4}\right)^0\right) - \left(\frac{1}{1-\frac{3}{4}} - \left(\frac{3}{4}\right)^0\right) = 5\left(\frac{4}{3}-1\right) - (4-1) = -\frac{4}{3}$

(b) $\displaystyle\sum_{n=1}^{\infty} \frac{5^n-3}{4^n}$ diverges because $\displaystyle\lim_{n\to\infty} \frac{5^n-3}{4^n} = \infty$ (instead of 0).

Problem 2. (4 points) Determine the following limits or state that the limit does not exist.

(a) $\displaystyle\lim_{n\to\infty} \frac{2^n-4^n}{7n^5+4^n} =$ -1

(b) $\displaystyle\lim_{n\to\infty} \frac{5n^2-n^3}{2n^2+4} =$ ∞

(c) $\displaystyle\lim_{n\to\infty} \sqrt{\frac{4n^3+1}{3+n+2n^3}} =$ $\sqrt{2}$

(d) $\displaystyle\lim_{n\to\infty} e^{1/n^2} =$ $e^0 = 1$

Armin Straub
straub@southalabama.edu

Problem 3. (10 points) Determine whether the following series converge or diverge. Make sure to indicate a reason!

(a) $\displaystyle\sum_{n=1}^{\infty} \frac{5n+1}{2n^2+1}$

☐ series converges ☐ series diverges

Indicate a reason:

The series diverges by limit comparison with the diverging harmonic series $\displaystyle\sum_{n=2}^{\infty} \frac{5n}{2n^2} = \frac{5}{2}\sum_{n=2}^{\infty} \frac{1}{n}$.

(b) $\displaystyle\sum_{n=2}^{\infty} \frac{4+3\log(n)}{2+\log(n)}$

☐ series converges ☐ series diverges

Indicate a reason:

The series diverges because $\dfrac{4+3\log(n)}{2+\log(n)} \to 3 \neq 0$ as $n \to \infty$.

(c) $\displaystyle\sum_{n=2}^{\infty} \frac{4^n}{n3^n}$

☐ series converges ☐ series diverges

Indicate a reason:

The series diverges because $\dfrac{4^n}{n3^n} \to \infty \neq 0$ as $n \to \infty$.

(d) $\displaystyle\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{3n+1}}$

☐ series converges ☐ series diverges

Indicate a reason:

The series converges by the alternating series test because $1/\sqrt{3n+1}$ is positive and decreases to 0.

(e) $\displaystyle\sum_{n=2}^{\infty} \frac{\sqrt{n}+2n+3}{4n^3+1}$

☐ series converges ☐ series diverges

Indicate a reason:

The series converges by limit comparison with the converging p-series $\displaystyle\sum_{n=1}^{\infty} \frac{2n}{4n^3} = \frac{1}{2}\sum_{n=1}^{\infty} \frac{1}{n^2}$ (because $p = 2 > 1$).

Problem 4. (10 points) Consider the power series $\displaystyle\sum_{n=1}^{\infty} \frac{(x+1)^n}{\sqrt{n}\,4^n}$.

(a) Determine the radius of convergence R.

(b) What is the exact interval of convergence?

(c) Let $f(x) = \displaystyle\sum_{n=1}^{\infty} \frac{(x+1)^n}{\sqrt{n}\,4^n}$ for x such that $|x+1| < R$. Write down a power series for $f'(x)$.

Solution.

(a) We apply the ratio test with $a_n = \dfrac{(x+1)^n}{\sqrt{n}\,4^n}$.

$$\left|\frac{a_{n+1}}{a_n}\right| = \left|\frac{(x+1)^{n+1}}{\sqrt{n+1}\,4^{n+1}} \frac{\sqrt{n}\,4^n}{(x+1)^n}\right| = \left|\frac{x+1}{4}\right| \frac{\sqrt{n}}{\sqrt{n+1}} \to \left|\frac{x+1}{4}\right| \text{ as } n \to \infty$$

Hence, the ratio test implies that $\displaystyle\sum_{n=1}^{\infty} \frac{(x+1)^n}{\sqrt{n}\,4^n}$ converges if $\left|\dfrac{x+1}{4}\right| < 1$, that is, $|x+1| < 4$.

In particular, the radius of convergence is 4.

(b) We already know that the series converges if $|x+1| < 4$, that is, if $x \in (-1-4, -1+4) = (-5, 3)$. We also know that the series diverges if $|x+1| > 4$. What we don't know yet is whether the series converges at the endpoints of the interval.

- $x = 3$: in that case, the series is $\displaystyle\sum_{n=1}^{\infty} \frac{(3+1)^n}{\sqrt{n}\,4^n} = \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$. This is the p-series with $p = \frac{1}{2}$ which we know diverges (because $p = \frac{1}{2} \leqslant 1$).

- $x = -5$: in that case, the series is $\displaystyle\sum_{n=1}^{\infty} \frac{(-5+1)^n}{\sqrt{n}\,4^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$. This is the alternating p-series with $p = \frac{1}{2}$ which we know converges (because $p = \frac{1}{2} > 0$).

  (We can use the alternating series test if we don't recall this fact about alternating p-series.)

In conclusion, the exact interval of convergence is $[-5, 3)$.

(c) $\displaystyle f'(x) = \sum_{n=1}^{\infty} \frac{n(x+1)^{n-1}}{\sqrt{n}\,4^n} = \sum_{n=1}^{\infty} \frac{\sqrt{n}(x+1)^{n-1}}{4^n}$

Optional: By replacing the index n with $n+1$, we can rewrite this as $\displaystyle f'(x) = \sum_{n=0}^{\infty} \frac{\sqrt{n+1}(x+1)^n}{4^{n+1}}$.

(extra scratch paper)