

**Review.** elliptic curves, Birch–Swinnerton-Dyer (BSD) conjecture, a few words about Bitcoin

**Example 200. (Riemann hypothesis)** The **Riemann zeta function**  $\zeta(s) = \sum_{n \geq 1} \frac{1}{n^s}$  converges (for real  $s$ ) if and only if  $s > 1$ .

The divergent series  $\zeta(1)$  is the harmonic series, and  $\zeta(p)$  is often called a  $p$ -series in Calculus II.

**Comment.** Euler achieved worldwide fame by discovering and proving that  $\zeta(2) = \frac{\pi^2}{6}$  (and similar formulas for  $\zeta(4), \zeta(6), \dots$ ).

For complex values of  $s \neq 1$ , there is a unique way to “analytically continue” this function. It is then “easy” to see that  $\zeta(-2) = 0, \zeta(-4) = 0, \dots$ . The **Riemann hypothesis** claims that all other zeroes of  $\zeta(s)$  lie on the line  $s = \frac{1}{2} + a\sqrt{-1}$  ( $a \in \mathbb{R}$ ). A proof of this conjecture (checked for the first 10,000,000,000 zeroes) is worth \$1,000,000.

<http://www.claymath.org/millennium-problems/riemann-hypothesis>

**The connection to primes.** Here’s a vague indication that  $\zeta(s)$  is intimately connected to prime numbers:

$$\begin{aligned} \zeta(s) &= \left(1 + \frac{1}{2^s} + \frac{1}{2^{2s}} + \dots\right) \left(1 + \frac{1}{3^s} + \frac{1}{3^{2s}} + \dots\right) \left(1 + \frac{1}{5^s} + \frac{1}{5^{2s}} + \dots\right) \dots \\ &= \frac{1}{1-2^{-s}} \frac{1}{1-3^{-s}} \frac{1}{1-5^{-s}} \dots \\ &= \prod_{p \text{ prime}} \frac{1}{1-p^{-s}} \end{aligned}$$

This infinite product is called the Euler product for the zeta function. If the Riemann hypothesis was true, then we would be better able to estimate the number  $\pi(x)$  of primes  $p \leq x$ .

More generally, certain statements about the zeta function can be translated to statements about primes. For instance, the (non-obvious!) fact that  $\zeta(s)$  has no zeros for  $\text{Re } s = 1$  implies the prime number theorem.

<http://www-users.math.umn.edu/~garrett/m/v/pnt.pdf>