

Example 28. (bonus challenge!) Eve, can you crack the following message?

KOBOSNOXSGYOM

Word on the street is that Alice was using the Vigenere cipher with a key of size 2.

(To collect a bonus point, send me an email before next class with the plaintext and how you found it.)

Attacks

So far, we considered the weakest kind of attack only: namely, a **ciphertext only attack**. And, even then, the historical ciphers prove to be terribly insecure.

However, we need to also worry about attacks where our enemy has additional insight.

- In a **known plaintext attack**, the enemy somehow has knowledge of a plaintext-ciphertext pair (m, c) .
- In a **chosen plaintext attack**, the enemy can, herself, compute $c = E(m)$ for a chosen plaintext m (“gained some sort of access to our encryption device”).
- In a **chosen ciphertext attack**, the enemy can, herself, compute $m = D(c)$ for a chosen ciphertext c (“gained some sort of access to our decryption device”).

There exist many variations of these. Sometimes, the attacker can make several choices (maybe even adaptively), sometimes she only has partial information.

Example 29. Alice sends the ciphertext *BKNDKGBQ* to Bob. Somehow, Eve has learned that Alice is using the Vigenere cipher and that the plaintext is *ALLCLEAR*. Next day, Alice sends the message *DNFFQGE*. Crack it and figure out the key that Alice used! (What kind of attack is this?)

Solution. This is a known plaintext attack.

Since $m + k = c$ (to be interpreted characterwise, modulo 26, and with k repeated as necessary), we can find k simply as $k = c - m$.

For instance, since *A* (value 0!) got encrypted to *B*, the first letter of the key is *B*.

<i>c</i>		<i>B</i>	<i>K</i>	<i>N</i>	<i>D</i>	<i>K</i>	<i>G</i>	<i>B</i>	<i>Q</i>
<i>m</i>	-	<i>A</i>	<i>L</i>	<i>L</i>	<i>C</i>	<i>L</i>	<i>E</i>	<i>A</i>	<i>R</i>
<i>k</i>	=	<i>B</i>	<i>Z</i>	<i>C</i>	<i>B</i>	<i>Z</i>	<i>C</i>	<i>B</i>	<i>Z</i>

We conclude that the key is $k = BZC$.

Note. Now, we can decrypt any future message that Alice sends using this key. For instance, we easily decrypt *DNFFQGE* to *CODERED* (using $m = c - k$).

All of the historical ciphers we have seen, including the substitution cipher that we will discuss shortly, fall apart completely under a known plaintext attack.

Euler's theorem

Example 30. Compute $3^{1003} \pmod{101}$.

Solution. Since 101 is a prime, $3^{100} \equiv 1 \pmod{101}$ by Fermat's little theorem. Because $3^{100} \equiv 3^0 \pmod{101}$, this enables us to reduce exponents modulo 100. In particular, since $1003 \equiv 3 \pmod{100}$, we have $3^{1003} \equiv 3^3 = 27 \pmod{101}$.

Fermat's little theorem is a special case of Euler's theorem :

Theorem 31. (Euler's theorem) If $n \geq 1$ and $\gcd(a, n) = 1$, then $a^{\phi(n)} \equiv 1 \pmod{n}$.

Proof. Euler's theorem can be proved along the lines of our earlier proof of Fermat's little theorem. The only adjustment is to only start with multiples ka where k is invertible modulo n . There are $\phi(n)$ such residues k , and so that's where Euler's phi function comes in. Can you complete the proof? \square

Example 32. What are the last two (decimal) digits of 3^{7082} ?

Solution. We need to determine $3^{7082} \pmod{100}$. $\phi(100) = \phi(2^2 5^2) = \phi(2^2)\phi(5^2) = (2^2 - 2^1)(5^2 - 5^1) = 40$. Since $\gcd(3, 100) = 1$ and $7082 \equiv 2 \pmod{40}$, Euler's theorem shows that $3^{7082} \equiv 3^2 = 9 \pmod{100}$.

Binary exponentiation

Example 33. Compute $3^{25} \pmod{101}$.

Solution. Fermat's little theorem is not helpful here.

Instead, we do **binary exponentiation**:

$3^2 = 9$, $3^4 = 81 \equiv -20$, $3^8 \equiv (-20)^2 = 400 \equiv -4$, $3^{16} \equiv (-4)^2 \equiv 16$, all modulo 101

$25 = 16 + 8 + 1$ [Every integer $n \geq 0$ can be written as a sum of distinct powers of 2 (in a unique way).]

Hence, $3^{25} = 3^{16} \cdot 3^8 \cdot 3^1 \equiv 16 \cdot (-4) \cdot 3 = -192 \equiv 10 \pmod{101}$.

Example 34. (extra practice) Compute $2^{20} \pmod{41}$.

Solution. $2^2 = 4$, $2^4 = 16$, $2^8 = 256 \equiv 10$, $2^{16} \equiv 100 \equiv 18$. Hence, $2^{20} = 2^{16} \cdot 2^4 \equiv 18 \cdot 16 = 288 \equiv 1 \pmod{41}$.

Or: $2^5 = 32 \equiv -9 \pmod{41}$. Hence, $2^{20} = (2^5)^4 \equiv (-9)^4 = 81^2 \equiv (-1)^2 = 1 \pmod{41}$.

Comment. Write $a = 2^{20} \pmod{41}$. It follows from Fermat's little theorem that $a^2 = 2^{40} \equiv 1 \pmod{41}$. The argument below shows that $a \equiv \pm 1 \pmod{41}$ [but we don't know which until we do the calculation].

The equation $x^2 \equiv 1 \pmod{p}$ is equivalent to $(x-1)(x+1) \equiv 0 \pmod{p}$ [b/c $(x-1)(x+1) = x^2 - 1$]. Since p is a prime and $p \nmid (x-1)(x+1)$, we must have $p \mid (x-1)$ or $p \mid (x+1)$. In other words, $x \equiv \pm 1 \pmod{p}$.