

Example 78. (review) Determine the general solution of $y'' + 4y = 12x$.

Solution. The DE is $p(D)y = 12x$ with $p(D) = D^2 + 4$, which has roots $\pm 2i$. Thus, the general solution is $y(x) = y_p(x) + C_1\cos(2x) + C_2\sin(2x)$. It remains to find a particular solution y_p .

Since $D^2 \cdot (12x) = 0$, we apply D^2 to both sides of the DE to get the homogeneous DE $D^2(D^2 + 4) \cdot y = 0$. Its general solution is $C_1 + C_2x + C_3\cos(2x) + C_4\sin(2x)$ and y_p must be of this form. Indeed, there must be a particular solution of the simpler form $y_p = C_1 + C_2x$ (because $C_3\cos(2x) + C_4\sin(2x)$ can be added to any y_p). It remains to find appropriate values C_1, C_2 such that $y_p'' + 4y_p = 12x$. Since $y_p'' + 4y_p = 4C_1 + 4C_2x$, comparing coefficients yields $4C_1 = 0$ and $4C_2 = 12$, so that $C_1 = 0$ and $C_2 = 3$. In other words, $y_p = 3x$.

Therefore, the general solution to the original DE is $y(x) = 3x + C_1\cos(2x) + C_2\sin(2x)$.

Example 79. Determine the general solution of $y'' + 4y' + 4y = e^{3x}$.

Solution. The DE is $p(D)y = e^{3x}$ with $p(D) = D^2 + 4D + 4 = (D + 2)^2$, which has roots $-2, -2$. Thus, the general solution is $y(x) = y_p(x) + (C_1 + C_2x)e^{-2x}$. It remains to find a particular solution y_p .

Since $(D - 3)e^{3x} = 0$, we apply $(D - 3)$ to the DE to get the homogeneous DE $(D - 3)(D + 2)^2y = 0$.

Its general solution is $(C_1 + C_2x)e^{-2x} + C_3e^{3x}$ and y_p must be of this form. Indeed, there must be a particular solution of the simpler form $y_p = Ce^{3x}$.

To determine the value of C , we plug into the original DE: $y_p'' + 4y_p' + 4y_p = (9 + 4 \cdot 3 + 4)Ce^{3x} = e^{3x}$. Hence, $C = 1/25$. Therefore, the general solution to the original DE is $y(x) = (C_1 + C_2x)e^{-2x} + \frac{1}{25}e^{3x}$.

We found a recipe for solving nonhomogeneous linear DEs with constant coefficients.

Our approach works for $p(D)y = f(x)$ whenever the right-hand side $f(x)$ is the solution of some homogeneous linear DE with constant coefficients: $q(D)f(x) = 0$

Theorem 80. (method of undetermined coefficients) To find a particular solution y_p to an inhomogeneous linear DE with constant coefficients $p(D)y = f(x)$:

- Find $q(D)$ so that $q(D)f(x) = 0$. [This does not work for all $f(x)$.]
- Let r_1, \dots, r_n be the (“old”) roots of the polynomial $p(D)$.
Let s_1, \dots, s_m be the (“new”) roots of the polynomial $q(D)$.
- It follows that y_p solves the homogeneous DE $q(D)p(D)y = 0$.

The characteristic polynomial of this DE has roots $r_1, \dots, r_n, s_1, \dots, s_m$.

Let v_1, \dots, v_m be the “new” solutions (i.e. not solutions of the “old” $p(D)y = 0$).

By plugging into $p(D)y_p = f(x)$, we find (unique) C_i so that $y_p = C_1v_1 + \dots + C_mv_m$.

Because of the final step, this approach is often called **method of undetermined coefficients**.

For which $f(x)$ does this work? By Theorem 64, we know exactly which $f(x)$ are solutions to homogeneous linear DEs with constant coefficients: these are linear combinations of exponentials $x^j e^{rx}$ (which includes $x^j e^{ax} \cos(bx)$ and $x^j e^{ax} \sin(bx)$).

Example 81. (again) Determine the general solution of $y'' + 4y' + 4y = e^{3x}$.

Solution. The “old” roots are $-2, -2$. The “new” roots are 3 . Hence, there has to be a particular solution of the form $y_p = Ce^{3x}$. To find the value of C , we plug into the DE.

$$y_p'' + 4y_p' + 4y_p = (9 + 4 \cdot 3 + 4)Ce^{3x} = e^{3x}. \text{ Hence, } C = 1/25.$$

Therefore, the general solution is $y(x) = (C_1 + C_2x)e^{-2x} + \frac{1}{25}e^{3x}$.

Example 82. Determine the general solution of $y'' + 4y' + 4y = 7e^{-2x}$.

Solution. The “old” roots are $-2, -2$. The “new” roots are -2 . Hence, there has to be a particular solution of the form $y_p = Cx^2e^{-2x}$. To find the value of C , we plug into the DE.

$$y_p' = C(-2x^2 + 2x)e^{-2x}$$

$$y_p'' = C(4x^2 - 8x + 2)e^{-2x}$$

$$y_p'' + 4y_p' + 4y_p = 2Ce^{-2x} = 7e^{-2x}$$

It follows that $C = 7/2$, so that $y_p = \frac{7}{2}x^2e^{-2x}$. The general solution is $y(x) = \left(C_1 + C_2x + \frac{7}{2}x^2\right)e^{-2x}$.