

**Review.** A homogeneous linear DE with constant coefficients is of the form  $p(D)y = 0$ , where  $p(D)$  is the characteristic polynomial. For each characteristic root  $r$  of multiplicity  $k$ , we get the  $k$  solutions  $x^j e^{rx}$  for  $j = 0, 1, \dots, k - 1$ .

**Example 75. (review)** Find the general solution of  $y''' + 2y'' + y' = 0$ .

**Solution.** The characteristic polynomial  $p(D) = D(D+1)^2$  has roots 0, 1, 1.  
Hence, the general solution is  $A + (B + Cx)e^x$ .

**Example 76.** Determine the general solution of  $y''' - 3y'' + 3y' - y = 0$ .

**Solution.** The characteristic polynomial  $p(D) = D^3 - 3D^2 + 3D - 1 = (D-1)^3$  has roots 1, 1, 1.  
By Theorem 72, the general solution is  $y(x) = (C_1 + C_2x + C_3x^2)e^x$ .

**Comment.** The coefficients 1, 2, 1 and 1, 3, 3, 1 in  $(D+1)^2$  and  $(D+1)^3$  are known as binomial coefficients. They can be arranged as rows in Pascal's triangle where the next row would be 1, 4, 6, 4, 1.

**Example 77.** Determine the general solution of  $y''' - y'' - 5y' - 3y = 0$ .

**Solution.** The characteristic polynomial  $p(D) = D^3 - D^2 - 5D - 3 = (D-3)(D+1)^2$  has roots 3, -1, -1.  
Hence, the general solution is  $y(x) = C_1 e^{3x} + (C_2 + C_3x)e^{-x}$ .

**Example 78. (homework)** Solve the IVP  $y''' = 8y'' - 16y'$  with  $y(0) = 1$ ,  $y'(0) = 4$ ,  $y''(0) = 0$ .

**Solution.** The characteristic polynomial  $p(D) = D^3 - 8D^2 + 16D = D(D-4)^2$  has roots 0, 4, 4.

By Theorem 72, the general solution is  $y(x) = C_1 + (C_2 + C_3x)e^{4x}$ .

Using  $y'(x) = (4C_2 + C_3 + 4C_3x)e^{4x}$  and  $y''(x) = 4(4C_2 + 2C_3 + 4C_3x)e^{4x}$ , the initial conditions result in the equations  $C_1 + C_2 = 1$ ,  $4C_2 + C_3 = 4$ ,  $16C_2 + 8C_3 = 0$ .

Solving these (start with the last two equations) we find  $C_1 = -1$ ,  $C_2 = 2$ ,  $C_3 = -4$ .

Hence the unique solution to the IVP is  $y(x) = -1 + (2 - 4x)e^{4x}$ .

**Important comment.** Check that  $y(x)$  indeed solves the IVP.

**Example 79.** Determine the general solution of  $y^{(6)} = 3y^{(5)} - 4y'''$ .

**Solution.** This DE is of the form  $p(D)y = 0$  with  $p(D) = D^6 - 3D^5 + 4D^3 = D^3(D-2)^2(D+1)$ .

The characteristic roots are 2, 2, 0, 0, 0, -1.

By Theorem 72, the general solution is  $y(x) = (C_1 + C_2x)e^{2x} + C_3 + C_4x + C_5x^2 + C_6e^{-x}$ .

**Example 80.** Consider the function  $y(x) = 3xe^{-2x} + 7e^x$ . Determine a homogeneous linear DE with constant coefficients of which  $y(x)$  is a solution.

**Solution.** In order for  $y(x)$  to be a solution of  $p(D)y = 0$ , the characteristic roots must include -2, -2, 1.

The simplest choice for  $p(D)$  thus is  $p(D) = (D+2)^2(D-1) = D^3 + 3D^2 - 4$ .

Accordingly,  $y(x)$  is a solution of  $y''' + 3y'' - 4y = 0$ .

**Example 81.** Consider the function  $y(x) = 3xe^{-2x} + 7$ . Determine a homogeneous linear DE with constant coefficients of which  $y(x)$  is a solution.

**Solution.** In order for  $y(x)$  to be a solution of  $p(D)y = 0$ , the characteristic roots must include -2, -2, 0.

The simplest choice for  $p(D)$  thus is  $p(D) = (D+2)^2D = D^3 + 4D^2 + 4D$ .

Accordingly,  $y(x)$  is a solution of  $y''' + 4y'' + 4y' = 0$ .

## Real form of complex solutions

Let's recall some basic facts about **complex numbers**:

- Every complex number can be written as  $z = x + iy$  with real  $x, y$ .
- Here, the imaginary unit  $i$  is characterized by solving  $x^2 = -1$ .  
**Important observation.** The same equation is solved by  $-i$ . This means that, algebraically, we cannot distinguish between  $+i$  and  $-i$ .
- The **conjugate** of  $z = x + iy$  is  $\bar{z} = x - iy$ .  
**Important comment.** Since we cannot algebraically distinguish between  $\pm i$ , we also cannot distinguish between  $z$  and  $\bar{z}$ . That's the reason why, in problems involving only real numbers, if a complex number  $z = x + iy$  shows up, then its **conjugate**  $\bar{z} = x - iy$  has to show up in the same manner. With that in mind, have another look at the examples below.
- The **real part** of  $z = x + iy$  is  $x$  and we write  $\operatorname{Re}(z) = x$ .  
Likewise the **imaginary part** is  $\operatorname{Im}(z) = y$ .  
Observe that  $\operatorname{Re}(z) = \frac{1}{2}(z + \bar{z})$  as well as  $\operatorname{Im}(z) = \frac{1}{2i}(z - \bar{z})$ .

### Theorem 82. (Euler's identity) $e^{ix} = \cos(x) + i \sin(x)$

**Proof.** Observe that both sides are the (unique) solution to the IVP  $y' = iy, y(0) = 1$ .

[Check that by computing the derivatives and verifying the initial condition! As we did in class.]  $\square$

**On lots of T-shirts.** In particular, with  $x = \pi$ , we get  $e^{\pi i} = -1$  or  $e^{i\pi} + 1 = 0$  (which connects the five fundamental constants).

**Comment.** It follows that  $\cos(x) = \operatorname{Re}(e^{ix}) = \frac{1}{2}(e^{ix} + e^{-ix})$  and  $\sin(x) = \operatorname{Im}(e^{ix}) = \frac{1}{2i}(e^{ix} - e^{-ix})$ .

**Example 83.** Determine the general solution of  $y'' + y = 0$ .

**Solution. (complex numbers in general solution)** The characteristic polynomial is  $D^2 + 1$  which has no roots over the reals. Over the **complex numbers**, by definition, the roots are  $i$  and  $-i$ .

So the general solution is  $y(x) = C_1 e^{ix} + C_2 e^{-ix}$ .

**Solution. (real general solution)** On the other hand, we easily check that  $y_1 = \cos(x)$  and  $y_2 = \sin(x)$  are two solutions. Hence, the general solution can also be written as  $y(x) = D_1 \cos(x) + D_2 \sin(x)$ .

**Important comment.** That we have these two different representations is a consequence of Euler's identity (Theorem 82) by which  $e^{\pm ix} = \cos(x) \pm i \sin(x)$ .

On the other hand,  $\cos(x) = \frac{1}{2}(e^{ix} + e^{-ix})$  and  $\sin(x) = \frac{1}{2i}(e^{ix} - e^{-ix})$ .

[Recall that the first formula is an instance of  $\operatorname{Re}(z) = \frac{1}{2}(z + \bar{z})$  and the second of  $\operatorname{Im}(z) = \frac{1}{2i}(z - \bar{z})$ .]

**Example 84.** Determine the general solution of  $y'' - 4y' + 13y = 0$  using only real numbers.

**Solution.** The characteristic polynomial  $p(D) = D^2 - 4D + 13$  has roots  $2 + 3i, 2 - 3i$ .

[We can use the quadratic formula to find these roots as  $\frac{4 \pm \sqrt{4^2 - 4 \cdot 13}}{2} = \frac{4 \pm \sqrt{-36}}{2} = \frac{4 \pm 6i}{2} = 2 \pm 3i$ .]

Hence, the general solution in real form is  $y(x) = C_1 e^{2x} \cos(3x) + C_2 e^{2x} \sin(3x)$ .

**Note.**  $e^{(2 \pm 3i)x} = e^{2x} e^{\pm 3ix} = e^{2x} (\cos(3x) \pm i \sin(3x))$

**Review.** A linear DE of order  $n$  is of the form

$$y^{(n)} + P_{n-1}(x)y^{(n-1)} + \dots + P_1(x)y' + P_0(x)y = f(x).$$

The **general solution of linear DE** always takes the form

$$y(x) = y_p(x) + C_1y_1(x) + \dots + C_ny_n(x),$$

where  $y_p$  is any solution (called a **particular solution**) and  $y_1, y_2, \dots, y_n$  are solutions to the corresponding **homogeneous** linear DE.

- In terms of  $D = \frac{d}{dx}$ , the DE becomes:  $Ly = f(x)$  with  $L = D^n + P_{n-1}(x)D^{n-1} + \dots + P_1(x)D + P_0(x)$ .
- The inclusion of the  $f(x)$  term makes  $Ly = f(x)$  an **inhomogeneous** linear DE. The corresponding **homogeneous** DE is  $Ly = 0$  (note that the zero function  $y(x) = 0$  is a solution of  $Ly = 0$ ).
- $L$  is called a **linear differential operator**.
  - $L(C_1y_1 + C_2y_2) = C_1Ly_1 + C_2Ly_2$  (**linearity**)  
**Comment.** If you are familiar with linear algebra, think of  $L$  replaced with a matrix  $A$  and  $y_1, y_2$  replaced with vectors  $v_1, v_2$ . In that case, the same linearity property holds.
  - So, if  $y_1$  solves  $Ly = f(x)$ , and  $y_2$  solves  $Ly = g(x)$ , then  $C_1y_1 + C_2y_2$  solves  $C_1f(x) + C_2g(x)$ .
  - In particular, if  $y_1$  and  $y_2$  solve the homogeneous DE, then so does any linear combination  $C_1y_1 + C_2y_2$ . This explains why, for any homogeneous linear DE of order  $n$ , there are  $n$  solutions  $y_1, y_2, \dots, y_n$  such that the general solution is  $y(x) = C_1y_1(x) + \dots + C_ny_n(x)$ . Moreover, in that case, if we have a **particular solution**  $y_p$  of the inhomogeneous DE  $Ly = f(x)$ , then  $y_p + C_1y_1 + \dots + C_ny_n$  is the general solution of  $Ly = f(x)$ .

**Example 85. (preview)** Determine the general solution of  $y'' + 4y = 12x$ . *Hint:*  $3x$  is a solution.

**Solution.** Here,  $p(D) = D^2 + 4$ . Because of the hint, we know that a particular solution is  $y_p = 3x$ .

The homogeneous DE  $p(D)y = 0$  has solutions  $y_1 = \cos(2x)$  and  $y_2 = \sin(2x)$ . [Make sure this is clear!]

Therefore, the general solution to the original DE is  $y_p + C_1y_1 + C_2y_2 = 3x + C_1\cos(2x) + C_2\sin(2x)$ .

**Just to make sure.** The DE in operator notation is  $Ly = f(x)$  with  $L = D^2 + 4$  and  $f(x) = 12x$ .

**Next.** How to find the particular solution  $y_p = 3x$  ourselves.