

Systems of differential equations

Example 69. (review) Write the (second-order) differential equation $y'' = 2y' + y$ as a system of (first-order) differential equations.

Solution. If $\mathbf{y} = \begin{bmatrix} y \\ y' \end{bmatrix}$, then $\mathbf{y}' = \begin{bmatrix} y' \\ y'' \end{bmatrix} = \begin{bmatrix} y' \\ 2y' + y \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} y \\ y' \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 2 \end{bmatrix} \mathbf{y}$. For short, $\mathbf{y}' = \begin{bmatrix} 0 & 1 \\ 1 & 2 \end{bmatrix} \mathbf{y}$.

Comment. Hence, we care about systems of differential equations, even if we work with just one function.

Example 70. Write the (third-order) differential equation $y''' = 3y'' - 2y' + y$ as a system of (first-order) differential equations.

Solution. If $\mathbf{y} = \begin{bmatrix} y \\ y' \\ y'' \end{bmatrix}$, then $\mathbf{y}' = \begin{bmatrix} y' \\ y'' \\ y''' \end{bmatrix} = \begin{bmatrix} y' \\ y'' \\ 3y'' - 2y' + y \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -2 & 3 \end{bmatrix} \begin{bmatrix} y \\ y' \\ y'' \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -2 & 3 \end{bmatrix} \mathbf{y}$.

Example 71. Consider the following system of (second-order) initial value problems:

$$\begin{aligned} y_1'' &= 2y_1' - 3y_2' + 7y_2 & y_1(0) &= 2, \quad y_1'(0) = 3, \quad y_2(0) = -1, \quad y_2'(0) = 1 \\ y_2'' &= 4y_1' + y_2' - 5y_1 \end{aligned}$$

Write it as a first-order initial value problem in the form $\mathbf{y}' = M\mathbf{y}$, $\mathbf{y}(0) = \mathbf{y}_0$.

Solution. If $\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_1' \\ y_2' \end{bmatrix}$, then $\mathbf{y}' = \begin{bmatrix} y_1' \\ y_2' \\ y_1'' \\ y_2'' \end{bmatrix} = \begin{bmatrix} y_1' \\ y_2' \\ 2y_1' - 3y_2' + 7y_2 \\ 4y_1' + y_2' - 5y_1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 7 & 2 & -3 \\ -5 & 0 & 4 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_1' \\ y_2' \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 7 & 2 & -3 \\ -5 & 0 & 4 & 1 \end{bmatrix} \mathbf{y}$.

For short, the system translates into $\mathbf{y}' = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 7 & 2 & -3 \\ -5 & 0 & 4 & 1 \end{bmatrix} \mathbf{y}$ with $\mathbf{y}(0) = \begin{bmatrix} 2 \\ -1 \\ 3 \\ 1 \end{bmatrix}$.

To solve $\mathbf{y}' = M\mathbf{y}$, determine the eigenvectors of M .

- Each λ -eigenvector \mathbf{v} provides a solution: $\mathbf{y}(x) = \mathbf{v}e^{\lambda x}$
- If there are enough eigenvectors, these combine to the general solution.

(systems of DEs) The unique solution to $\mathbf{y}' = M\mathbf{y}$, $\mathbf{y}(0) = \mathbf{c}$ is $\mathbf{y}(x) = e^{Mx}\mathbf{c}$.

- Here, e^{Mx} is the fundamental matrix solution to $\mathbf{y}' = M\mathbf{y}$, $\mathbf{y}(0) = I$ (with I the identity matrix).
- If $\Phi(x)$ is any fundamental matrix solution to $\mathbf{y}' = M\mathbf{y}$, then $e^{Mx} = \Phi(x)\Phi(0)^{-1}$.
- To construct a fundamental matrix solution $\Phi(x)$, we compute eigenvectors:
Given a λ -eigenvector \mathbf{v} , we have the corresponding solution $\mathbf{y}(x) = \mathbf{v}e^{\lambda x}$.
If there are enough eigenvectors, we can collect these as columns to obtain $\Phi(x)$.

Note. We are defining the **matrix exponential** e^{Mx} as the solution to an IVP. This is equivalent to how one can define the ordinary exponential e^x as the solution to $y' = y$, $y(0) = 1$.

[In a little bit, we will also discuss how to think about the matrix exponential e^{Mx} using power series.]

Comment. If there are not enough eigenvectors, then we know what to do (at least in principle): instead of looking only for solutions of the type $\mathbf{y}(x) = \mathbf{v}e^{\lambda x}$, we also need to look for solutions of the type $\mathbf{y}(x) = (\mathbf{v}x + \mathbf{w})e^{\lambda x}$. Note that this can only happen if an eigenvalue is a repeated root of the characteristic polynomial.

Important. Compare this to our method of solving systems of REs and for computing matrix powers M^n . Note that the above conclusion about systems of DEs can be deduced along the same lines as what we did for REs:

- If $\Phi(x)$ is a fundamental matrix solution, then so is $\Psi(x) = \Phi(x)C$ for every constant matrix C . (Why?!) Therefore, $\Psi(x) = \Phi(x)\Phi(0)^{-1}$ is a fundamental matrix solution with $\Psi(0) = \Phi(0)\Phi(0)^{-1} = I$. But e^{Mx} is defined to be the unique such solution, so that $\Psi(x) = e^{Mx}$.
- Let us look for solutions of $\mathbf{y}' = M\mathbf{y}$ of the form $\mathbf{y}(x) = \mathbf{v}e^{\lambda x}$. Note that $\mathbf{y}' = \lambda \mathbf{v}e^{\lambda x} = \lambda \mathbf{y}$. Plugging into $\mathbf{y}' = M\mathbf{y}$, we find $\lambda \mathbf{y} = M\mathbf{y}$. In other words, $\mathbf{y}(x) = \mathbf{v}e^{\lambda x}$ is a solution if and only if \mathbf{v} is a λ -eigenvector of M .

Observe how the next example proceeds along the same lines as Example 62.

Important. In fact, we can translate back and forth (without additional computations) by simply replacing 3^n and $(-2)^n$ by e^{3x} and e^{-2x} .

Example 72. (homework) Let $M = \begin{bmatrix} 8 & -10 \\ 5 & -7 \end{bmatrix}$.

- Determine the general solution to $\mathbf{y}' = M\mathbf{y}$.
- Determine a fundamental matrix solution to $\mathbf{y}' = M\mathbf{y}$.
- Compute e^{Mx} .
- Solve the initial value problem $\mathbf{y}' = M\mathbf{y}$ with $\mathbf{y}(0) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Solution.

- Recall that each λ -eigenvector \mathbf{v} of M provides us with a solution: namely, $\mathbf{y}(x) = \mathbf{v}e^{\lambda x}$. We computed earlier that $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvector for $\lambda = 3$, and $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is an eigenvector for $\lambda = -2$. Hence, the general solution is $C_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{3x} + C_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{-2x}$.
- The corresponding fundamental matrix solution is $\Phi(x) = \begin{bmatrix} 2 \cdot e^{3x} & e^{-2x} \\ e^{3x} & e^{-2x} \end{bmatrix}$. [Note that our general solution is precisely $\Phi(x) \begin{bmatrix} C_1 \\ C_2 \end{bmatrix}$.]
- Since $\Phi(0) = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$, we have $\Phi(0)^{-1} = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$. It follows that

$$e^{Mx} = \Phi(x)\Phi(0)^{-1} = \begin{bmatrix} 2 \cdot e^{3x} & e^{-2x} \\ e^{3x} & e^{-2x} \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 2 \cdot e^{3x} - e^{-2x} & -2 \cdot e^{3x} + 2e^{-2x} \\ e^{3x} - e^{-2x} & -e^{3x} + 2e^{-2x} \end{bmatrix}.$$

Check. Let us verify the formula for e^{Mx} in the simple case $x = 0$: $e^{M0} = \begin{bmatrix} 2-1 & -2+2 \\ 1-1 & -1+2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

- The solution to the IVP is $\mathbf{y}(x) = e^{Mx} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \cdot e^{3x} + 2e^{-2x} \\ -e^{3x} + 2e^{-2x} \end{bmatrix}$ (the second column of e^{Mx}).

Sage. We can compute the matrix exponential in Sage as follows:

```
>>> M = matrix([[8,-10],[5,-7]])
```

```
>>> exp(M*x)
```

$$\begin{pmatrix} (2e^{5x} - 1)e^{-2x} & -2(e^{5x} - 1)e^{-2x} \\ (e^{5x} - 1)e^{-2x} & -(e^{5x} - 2)e^{-2x} \end{pmatrix}$$

Note that this indeed matches the result of our computation.

[By the way, the variable x is pre-defined as a symbolic variable in Sage. That's why, unlike for n in the computation of M^n , we did not need to use `x = var('x')` first.]

Example 73. Suppose that $e^{Mx} = \frac{1}{10} \begin{bmatrix} e^x + 9e^{2x} & 3e^x - 3e^{2x} \\ 3e^x - 3e^{2x} & 9e^x + e^{2x} \end{bmatrix}$.

- Without doing any computations, determine M^n .
- What is M ?
- Without doing any computations, determine the eigenvalues and eigenvectors of M .
- From these eigenvalues and eigenvectors, write down a simple fundamental matrix solution to $\mathbf{y}' = M\mathbf{y}$.
- From that fundamental matrix solution, how can we compute e^{Mx} ? (If we didn't know it already...)
- Having computed e^{Mx} , what is a simple check that we can (should!) make?

Solution.

- Since e^x and e^{2x} correspond to eigenvalues 1 and 2, we just need to replace these by $1^n = 1$ and 2^n :

$$M^n = \frac{1}{10} \begin{bmatrix} 1 + 9 \cdot 2^n & 3 - 3 \cdot 2^n \\ 3 - 3 \cdot 2^n & 9 + 2^n \end{bmatrix}$$

- We can simply set $n = 1$ in our formula for M^n , to get $M = \frac{1}{10} \begin{bmatrix} 19 & -3 \\ -3 & 11 \end{bmatrix}$.

- The eigenvalues are 1 and 2 (because e^{Mx} contains the exponentials e^x and e^{2x}).

Looking at the coefficients of e^x in the first column of e^{Mx} , we see that $\begin{bmatrix} 1 \\ 3 \end{bmatrix}$ is a 1-eigenvector.

[We can also look the second column of e^{Mx} , to obtain $\begin{bmatrix} 3 \\ 9 \end{bmatrix}$ which is a multiple and thus equivalent.]

Likewise, we find that $\begin{bmatrix} 9 \\ -3 \end{bmatrix}$ or, equivalently, $\begin{bmatrix} -3 \\ 1 \end{bmatrix}$ is a 2-eigenvector.

Comment. For the above to make sense, we need to keep in mind that, associated to a λ -eigenvector \mathbf{v} , we have the corresponding solution $\mathbf{y}(x) = \mathbf{v}e^{\lambda x}$ of the DE $\mathbf{y}' = M\mathbf{y}$. On the other hand, the columns of e^{Mx} are solutions to that DE and, therefore, must be linear combinations of these $\mathbf{v}e^{\lambda x}$.

- From the eigenvalues and eigenvectors, we know that $\begin{bmatrix} 1 \\ 3 \end{bmatrix}e^x$ and $\begin{bmatrix} -3 \\ 1 \end{bmatrix}e^{2x}$ are solutions (and that the general solutions consists of the linear combinations of these two).

Selecting these as the columns, we obtain the fundamental matrix solution $\Phi(x) = \begin{bmatrix} e^x & -3e^{2x} \\ 3e^x & e^{2x} \end{bmatrix}$.

Comment. The *fundamental* refers to the fact that the columns combine to the general solution.

The *matrix solution* means that $\Phi(x)$ itself satisfies the DE: namely, we have $\Phi' = M\Phi$. That this is the case is a consequence of matrix multiplication (namely, say, the second column of $M\Phi$ is defined to be M times the second column of Φ ; but that column is a vector solution and therefore solves the DE).

- We can compute e^{Mx} as $e^{Mx} = \Phi(x)\Phi(0)^{-1}$.

If $\Phi(x) = \begin{bmatrix} e^x & -3e^{2x} \\ 3e^x & e^{2x} \end{bmatrix}$, then $\Phi(0) = \begin{bmatrix} 1 & -3 \\ 3 & 1 \end{bmatrix}$ and, hence, $\Phi(0)^{-1} = \frac{1}{10} \begin{bmatrix} 1 & 3 \\ -3 & 1 \end{bmatrix}$. It follows that

$$e^{Mx} = \Phi(x)\Phi(0)^{-1} = \begin{bmatrix} e^x & -3e^{2x} \\ 3e^x & e^{2x} \end{bmatrix} \frac{1}{10} \begin{bmatrix} 1 & 3 \\ -3 & 1 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} e^x + 9e^{2x} & 3e^x - 3e^{2x} \\ 3e^x - 3e^{2x} & 9e^x + e^{2x} \end{bmatrix}.$$

- We can check that e^{Mx} equals the identity matrix if we set $x = 0$:

$$\frac{1}{10} \begin{bmatrix} e^x + 9e^{2x} & 3e^x - 3e^{2x} \\ 3e^x - 3e^{2x} & 9e^x + e^{2x} \end{bmatrix} \stackrel{x=0}{\rightsquigarrow} \frac{1}{10} \begin{bmatrix} 1+9 & 3-3 \\ 3-3 & 9+1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

This check does not require much effort and can even be done in our head while writing down e^{Mx} . There is really no excuse for not doing it!

Example 74. (homework) Let $M = \begin{bmatrix} -1 & 6 \\ -1 & 4 \end{bmatrix}$.

- Determine the general solution to $\mathbf{y}' = M\mathbf{y}$.
- Determine a fundamental matrix solution to $\mathbf{y}' = M\mathbf{y}$.
- Compute e^{Mx} .
- Solve the initial value problem $\mathbf{y}' = M\mathbf{y}$ with $\mathbf{y}(0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- Compute M^n .
- Solve $\mathbf{a}_{n+1} = M\mathbf{a}_n$ with $\mathbf{a}_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

Solution.

- (a) We determine the eigenvectors of M . The characteristic polynomial is:

$$\det(M - \lambda I) = \det\left(\begin{bmatrix} -1-\lambda & 6 \\ -1 & 4-\lambda \end{bmatrix}\right) = (-1-\lambda)(4-\lambda) + 6 = \lambda^2 - 3\lambda + 2 = (\lambda-1)(\lambda-2)$$

Hence, the eigenvalues are $\lambda = 1$ and $\lambda = 2$.

- $\lambda = 1$: Solving $\begin{bmatrix} -2 & 6 \\ -1 & 3 \end{bmatrix}\mathbf{v} = \mathbf{0}$, we find that $\mathbf{v} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$ is an eigenvector for $\lambda = 1$.
- $\lambda = 2$: Solving $\begin{bmatrix} -3 & 6 \\ -1 & 2 \end{bmatrix}\mathbf{v} = \mathbf{0}$, we find that $\mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvector for $\lambda = 2$.

Hence, the general solution is $C_1 \begin{bmatrix} 3 \\ 1 \end{bmatrix} e^x + C_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{2x}$.

- (b) The corresponding fundamental matrix solution is $\Phi = \begin{bmatrix} 3e^x & 2e^{2x} \\ e^x & e^{2x} \end{bmatrix}$.

- (c) Note that $\Phi(0) = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}$, so that $\Phi(0)^{-1} = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$. It follows that

$$e^{Mx} = \Phi(x)\Phi(0)^{-1} = \begin{bmatrix} 3e^x & 2e^{2x} \\ e^x & e^{2x} \end{bmatrix} \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix} = \begin{bmatrix} 3e^x - 2e^{2x} & -6e^x + 6e^{2x} \\ e^x - e^{2x} & -2e^x + 3e^{2x} \end{bmatrix}.$$

- (d) The solution to the IVP is $\mathbf{y}(x) = e^{Mx} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3e^x - 2e^{2x} & -6e^x + 6e^{2x} \\ e^x - e^{2x} & -2e^x + 3e^{2x} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -3e^x + 4e^{2x} \\ -e^x + 2e^{2x} \end{bmatrix}$.

Note. If we hadn't already computed e^{Mx} , we would use the general solution and solve for the appropriate values of C_1 and C_2 . Do it that way as well!

- (e) From the first part, it follows that $\mathbf{a}_{n+1} = M\mathbf{a}_n$ has general solution $C_1 \begin{bmatrix} 3 \\ 1 \end{bmatrix} + C_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} 2^n$.

(Note that $1^n = 1$.)

The corresponding fundamental matrix solution is $\Phi_n = \begin{bmatrix} 3 & 2 \cdot 2^n \\ 1 & 2^n \end{bmatrix}$.

As above, $\Phi_0 = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}$, so that $\Phi(0)^{-1} = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$ and

$$M^n = \Phi_n \Phi_0^{-1} = \begin{bmatrix} 3 & 2 \cdot 2^n \\ 1 & 2^n \end{bmatrix} \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix} = \begin{bmatrix} 3 - 2 \cdot 2^n & -6 + 6 \cdot 2^n \\ 1 - 2^n & -2 + 3 \cdot 2^n \end{bmatrix}.$$

Important. Compare with our computation for e^{Mx} . Can you see how this was basically the same computation? Write down M^n directly from e^{Mx} .

- (f) The (unique) solution is $\mathbf{a}_n = M^n \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 - 2 \cdot 2^n & -6 + 6 \cdot 2^n \\ 1 - 2^n & -2 + 3 \cdot 2^n \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -3 + 4 \cdot 2^n \\ -1 + 2 \cdot 2^n \end{bmatrix}$.

Again, compare with the earlier IVP!