

# Preparing for Midterm #2

Please print your name:

---

**Bonus challenge.** Let me know about any typos you spot in the posted solutions (or lecture sketches or any other posted material). Any typo, that is not yet fixed by the time you send it to me, is worth a bonus point.

**Problem 1.** Before working on the problems below, make sure that you have completed all homework. (In particular, Homework Sets 6, 7, 8 and 9.)

**Problem 2.** Solve the initial value problem  $\mathbf{y}' = \begin{bmatrix} 4 & -8 \\ -1 & 6 \end{bmatrix} \mathbf{y}$ ,  $\mathbf{y}(0) = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$ .

(Use Homework Problem 8.5 to generate more practice problems of this kind.)

**Solution.**

- $A = \begin{bmatrix} 4 & -8 \\ -1 & 6 \end{bmatrix}$  has characteristic polynomial  $(4 - \lambda)(6 - \lambda) - 8 = \lambda^2 - 10\lambda + 16 = (\lambda - 2)(\lambda - 8)$ .

Hence, the eigenvalues of  $A$  are 2, 8.

The 8-eigenspace  $\text{null}\left(\begin{bmatrix} -4 & -8 \\ -1 & -2 \end{bmatrix}\right)$  has basis  $\begin{bmatrix} -2 \\ 1 \end{bmatrix}$ .

The 2-eigenspace  $\text{null}\left(\begin{bmatrix} 2 & -8 \\ -1 & 4 \end{bmatrix}\right)$  has basis  $\begin{bmatrix} 4 \\ 1 \end{bmatrix}$ .

Hence,  $A = PDP^{-1}$  with  $P = \begin{bmatrix} -2 & 4 \\ 1 & 1 \end{bmatrix}$  and  $D = \begin{bmatrix} 8 & \\ & 2 \end{bmatrix}$ .

- Finally, we compute the solution  $\mathbf{y}(t) = e^{At}\mathbf{y}(0)$ :

$$\begin{aligned} \mathbf{y}(t) &= Pe^{Dt}P^{-1}\mathbf{y}(0) \\ &= \underbrace{\begin{bmatrix} -2 & 4 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} e^{8t} & \\ & e^{2t} \end{bmatrix} \begin{pmatrix} -\frac{1}{6} \end{pmatrix} \begin{bmatrix} 1 & -4 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix}}_{\begin{bmatrix} -2e^{8t} & 4e^{2t} \\ e^{8t} & e^{2t} \end{bmatrix}} \underbrace{\frac{1}{6} \begin{bmatrix} 1 \\ 5 \end{bmatrix}}_{\frac{1}{6} \begin{bmatrix} 1 \\ 5 \end{bmatrix}} = \frac{1}{6} \begin{bmatrix} 20e^{2t} - 2e^{8t} \\ 5e^{2t} + e^{8t} \end{bmatrix} \end{aligned}$$

**Problem 3.**

- (a) Convert the third-order differential equation

$$y''' = 6y'' - 3y' - 10y, \quad y(0) = 1, \quad y'(0) = 2, \quad y''(0) = 3$$

to a system of first-order differential equations.

- (b) Solve the original differential equation by solving the system.

(Use Homework Problems 8.6, 8.8 to generate more practice problems of this kind.)

**Solution.**

- (a) Write  $y_1 = y$ ,  $y_2 = y'$  and  $y_3 = y''$ .

Then,  $y''' = 6y'' - 3y' - 10y$  translates into the first-order system  $\begin{cases} y_1' = y_2 \\ y_2' = y_3 \\ y_3' = -10y_1 - 3y_2 + 6y_3 \end{cases}$ .

In matrix form, this is  $\mathbf{y}' = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -10 & -3 & 6 \end{bmatrix} \mathbf{y}$ ,  $\mathbf{y}(0) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ .

(b) First, to compute  $e^{At}$  for  $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -10 & -3 & 6 \end{bmatrix}$ , we need to diagonalize  $A$ .

- The eigenvalues of  $A$  are  $\lambda = 5, 2, -1$ .
- The 5-eigenspace  $\text{null}\left(\begin{bmatrix} -5 & 1 & 0 \\ 0 & -5 & 1 \\ -10 & -3 & 1 \end{bmatrix}\right)$  has basis  $\begin{bmatrix} 1 \\ 5 \\ 25 \end{bmatrix}$ .
- The 2-eigenspace  $\text{null}\left(\begin{bmatrix} -2 & 1 & 0 \\ 0 & -2 & 1 \\ -10 & -3 & 4 \end{bmatrix}\right)$  has basis  $\begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}$ .
- The  $-1$ -eigenspace  $\text{null}\left(\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ -10 & -3 & 7 \end{bmatrix}\right)$  has basis  $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$ .

Hence,  $A = PDP^{-1}$  with  $P = \begin{bmatrix} 1 & 1 & 1 \\ 5 & 2 & -1 \\ 25 & 4 & 1 \end{bmatrix}$  and  $D = \begin{bmatrix} 5 & & \\ & 2 & \\ & & -1 \end{bmatrix}$ .

Then, we compute the solution  $\mathbf{y} = e^{At}\mathbf{y}(0)$ :

$$\begin{aligned} \mathbf{y} = e^{At}\mathbf{y}(0) &= Pe^{Dt}P^{-1}\mathbf{y}(0) \\ &= \begin{bmatrix} 1 & 1 & 1 \\ 5 & 2 & -1 \\ 25 & 4 & 1 \end{bmatrix} \begin{bmatrix} e^{5t} & 0 & 0 \\ 0 & e^{2t} & 0 \\ 0 & 0 & e^{-t} \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 5 & 2 & -1 \\ 25 & 4 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 1 & 1 \\ 5 & 2 & -1 \\ 25 & 4 & 1 \end{bmatrix} \begin{bmatrix} e^{5t} & 0 & 0 \\ 0 & e^{2t} & 0 \\ 0 & 0 & e^{-t} \end{bmatrix} \frac{1}{18} \begin{bmatrix} -1 \\ 20 \\ -1 \end{bmatrix} \\ &= \frac{1}{18} \begin{bmatrix} 1 & 1 & 1 \\ 5 & 2 & -1 \\ 25 & 4 & 1 \end{bmatrix} \begin{bmatrix} -e^{5t} \\ 20e^{2t} \\ -e^{-t} \end{bmatrix} \\ &= \frac{1}{18} \begin{bmatrix} -e^{5t} + 20e^{2t} - e^{-t} \\ -5e^{5t} + 40e^{2t} + e^{-t} \\ -25e^{5t} + 80e^{2t} - e^{-t} \end{bmatrix} \end{aligned}$$

In particular, the original differential equation is solved by  $y(t) = \frac{1}{18}(-e^{5t} + 20e^{2t} - e^{-t})$ .

**Comment.** To compute  $\begin{bmatrix} 1 & 1 & 1 \\ 5 & 2 & -1 \\ 25 & 4 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \frac{1}{18} \begin{bmatrix} -1 \\ 20 \\ -1 \end{bmatrix}$ , we solve  $\begin{bmatrix} 1 & 1 & 1 \\ 5 & 2 & -1 \\ 25 & 4 & 1 \end{bmatrix} \mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  to find  $\mathbf{v} = \frac{1}{18} \begin{bmatrix} -1 \\ 20 \\ -1 \end{bmatrix}$ .

**Next comment.** Obviously, computations will be more pleasant on the exam.

#### Problem 4.

- What are the possible Jordan normal forms of a  $6 \times 6$  matrix with eigenvalues  $7, 7, 3, 3, 3, 3$ ?
- How many different Jordan normal forms are there for a  $10 \times 10$  matrix with eigenvalues  $8, 6, 6, 2, 2, 2, 1, 1, 1, 1$ ?

(Use Homework Problems 9.1, 9.2 to generate more practice problems of this kind.)

**Solution.**

(a) There are  $2 \cdot 5 = 10$  possibilities:

$$\begin{bmatrix} 7 & & & & \\ & 7 & & & \\ & & 3 & & \\ & & & 3 & \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & 1 \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & 1 \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & 1 \\ & & & & 3 \end{bmatrix},$$

$$\begin{bmatrix} 7 & 1 & & & \\ & 7 & & & \\ & & 3 & & \\ & & & 3 & \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & 1 & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & 1 & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & 1 \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & 1 & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & 1 \\ & & & & 3 \end{bmatrix}, \begin{bmatrix} 7 & 1 & & & \\ & 7 & & & \\ & & 3 & 1 & \\ & & & 3 & 1 \\ & & & & 3 \end{bmatrix}$$

(b) There are  $1 \cdot 2 \cdot 3 \cdot 5 = 30$  possible different Jordan normal forms.

**Problem 5.** Consider the sequence  $a_n$  defined by  $a_{n+2} = 4a_{n+1} - a_n$  and  $a_0 = 1, a_1 = 0$ .

(a) Determine the next three terms.

(b) A Binet-like formula for  $a_n$  is  $a_n =$  .

(c)  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} =$  .

(Use Homework Problems 7.1, 7.2, 7.3, 7.4, 7.5 to generate more practice problems of this kind.)

**Solution.**

(a)  $a_2 = -1, a_3 = -4, a_4 = -15$

(b) The recursion can be translated to  $\begin{bmatrix} a_{n+2} \\ a_{n+1} \end{bmatrix} = \begin{bmatrix} 4 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a_{n+1} \\ a_n \end{bmatrix}$ .

The eigenvalues of  $\begin{bmatrix} 4 & -1 \\ 1 & 0 \end{bmatrix}$  are  $2 + \sqrt{3} \approx 3.732$  and  $2 - \sqrt{3} \approx 0.268$ .

Hence,  $a_n = \alpha_1 (2 + \sqrt{3})^n + \alpha_2 (2 - \sqrt{3})^n$  and we only need to figure out the two unknowns  $\alpha_1, \alpha_2$ . We can do that using the two initial conditions:  $a_0 = \alpha_1 + \alpha_2 = 1, a_1 = (2 + \sqrt{3})\alpha_1 + (2 - \sqrt{3})\alpha_2 = 0$ .


Solving, we find  $\alpha_1 = \frac{\sqrt{3}-2}{2\sqrt{3}}$  and  $\alpha_2 = \frac{\sqrt{3}+2}{2\sqrt{3}}$  so that, in conclusion,  $a_n = \frac{\sqrt{3}-2}{2\sqrt{3}}(2 + \sqrt{3})^n + \frac{\sqrt{3}+2}{2\sqrt{3}}(2 - \sqrt{3})^n$ .

(c) It follows from the Binet-like formula that  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 2 + \sqrt{3}$ .

**Problem 6.** Suppose the internet consists of only the four webpages  $A, B, C, D$  which link to each other as indicated in the diagram.


Rank these webpages by computing their PageRank vector.

(Use Homework Problem 6.6 to generate more practice problems of this kind.)



**Solution.** Recall that we model a random surfer, who randomly clicks on links. Let  $a_t$  be the probability that such a surfer will be on page  $A$  at time  $t$ . Likewise,  $b_t, c_t, d_t$  are the probabilities that the surfer will be on page  $B, C$  or  $D$ .

The transition probabilities are indicated in the diagram to the right.



$$\begin{bmatrix} a_{t+1} \\ b_{t+1} \\ c_{t+1} \\ d_{t+1} \end{bmatrix} = \begin{bmatrix} 0 \cdot a_t + \frac{1}{2} \cdot b_t + \frac{1}{2} \cdot c_t + 1 \cdot d_t \\ \frac{1}{2} \cdot a_t + 0 \cdot b_t + 0 \cdot c_t + 0 \cdot d_t \\ \frac{1}{2} \cdot a_t + 0 \cdot b_t + 0 \cdot c_t + 0 \cdot d_t \\ 0 \cdot a_t + \frac{1}{2} \cdot b_t + \frac{1}{2} \cdot c_t + 0 \cdot d_t \end{bmatrix} = \underbrace{\begin{bmatrix} 0 & \frac{1}{2} & \frac{1}{2} & 1 \\ \frac{1}{2} & 0 & 0 & 0 \\ \frac{1}{2} & 0 & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 \end{bmatrix}}_{=T} \begin{bmatrix} a_t \\ b_t \\ c_t \\ d_t \end{bmatrix}$$

To find the equilibrium state, we determine an appropriate 1-eigenvector of the transition matrix  $T$ .

The 1-eigenspace is  $\text{null}(T - 1 \cdot I) = \text{null}\left(\begin{bmatrix} -1 & \frac{1}{2} & \frac{1}{2} & 1 \\ \frac{1}{2} & -1 & 0 & 0 \\ \frac{1}{2} & 0 & -1 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & -1 \end{bmatrix}\right)$

To compute a basis, we perform Gaussian elimination (details below):

$$\begin{bmatrix} -1 & \frac{1}{2} & \frac{1}{2} & 1 \\ \frac{1}{2} & -1 & 0 & 0 \\ \frac{1}{2} & 0 & -1 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

We conclude that the 1-eigenspace has basis  $\begin{bmatrix} 2 \\ 1 \\ 1 \\ 1 \end{bmatrix}$ . (Note that its entries add up to  $2 + 1 + 1 + 1 = 5$ .)

The corresponding equilibrium state is  $\frac{1}{5} \begin{bmatrix} 2 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0.4 \\ 0.2 \\ 0.2 \\ 0.2 \end{bmatrix}$ . This is the PageRank vector.

Correspondingly, we rank  $A$  the highest, followed by  $B, C, D$  which we rank equally.

[In hindsight, can you (at least sort of) see, directly from the diagram, why the PageRank is what it is?]

The full steps of the Gaussian elimination are:

$$\begin{bmatrix} -1 & \frac{1}{2} & \frac{1}{2} & 1 \\ \frac{1}{2} & -1 & 0 & 0 \\ \frac{1}{2} & 0 & -1 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & -1 \end{bmatrix} \xrightarrow{\substack{R_2 + \frac{1}{2}R_1 \Rightarrow R_2 \\ R_3 + \frac{1}{2}R_1 \Rightarrow R_3}} \begin{bmatrix} -1 & \frac{1}{2} & \frac{1}{2} & 1 \\ 0 & -\frac{3}{4} & \frac{1}{4} & \frac{1}{2} \\ 0 & \frac{1}{4} & -\frac{3}{4} & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} & -1 \end{bmatrix} \xrightarrow{\substack{R_3 + \frac{1}{3}R_2 \Rightarrow R_3 \\ R_4 + \frac{2}{3}R_2 \Rightarrow R_4}} \begin{bmatrix} -1 & \frac{1}{2} & \frac{1}{2} & 1 \\ 0 & -\frac{3}{4} & \frac{1}{4} & \frac{1}{2} \\ 0 & 0 & -\frac{2}{3} & \frac{2}{3} \\ 0 & 0 & \frac{2}{3} & -\frac{2}{3} \end{bmatrix} \xrightarrow{R_4 + R_3 \Rightarrow R_4} \begin{bmatrix} -1 & \frac{1}{2} & \frac{1}{2} & 1 \\ 0 & -\frac{3}{4} & \frac{1}{4} & \frac{1}{2} \\ 0 & 0 & -\frac{2}{3} & \frac{2}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\xrightarrow{\substack{-1R_1 \Rightarrow R_1 \\ -\frac{4}{3}R_2 \Rightarrow R_2 \\ -\frac{3}{2}R_3 \Rightarrow R_3}} \begin{bmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} & -1 \\ 0 & 1 & -\frac{1}{3} & -\frac{2}{3} \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{\substack{R_1 + \frac{1}{2}R_3 \Rightarrow R_1 \\ R_2 + \frac{1}{3}R_3 \Rightarrow R_2}} \begin{bmatrix} 1 & -\frac{1}{2} & 0 & -\frac{3}{2} \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 + \frac{1}{2}R_2 \Rightarrow R_1} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

This was good practice of elimination! However, notice that we can actually find an eigenvector  $\mathbf{x}$  with less effort by spelling out the equations: for instance, the second one is just  $\frac{1}{2}x_1 - x_2 = 0$ . Do that!

**Problem 7.** Determine an orthogonal matrix  $P$  and a diagonal matrix  $D$  such that  $A = PDP^T$ .

(a) If  $A$  is the  $3 \times 3$  matrix for reflecting through the plane spanned by the vectors  $\begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ .

(b) If  $A$  is the  $3 \times 3$  matrix for reflecting through the plane spanned by the vectors  $\begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ .

(Use Homework Problem 8.2 to generate more practice problems of this kind.)

**Solution.** In each case, let  $W$  be the plane spanned by the given two vectors. Then  $A$  has 1-eigenspace  $W$  and  $-1$ -eigenspace  $W^\perp$ . We need orthonormal bases for  $W$  and  $W^\perp$  in order to write down the diagonalization  $A = PDP^T$ .

**Important comment.** Note that, if we just use any bases for  $W$  and  $W^\perp$ , then we would only get a diagonalization of the type  $A = PDP^{-1}$ .

(a) Here,  $W = \text{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ .

- (basis for  $W$ ) Since the vectors are already orthogonal, we normalize to find that  $\frac{1}{\sqrt{5}} \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$  is an orthonormal basis for  $W$ .

- (basis for  $W^\perp$ ) We can read off that  $\frac{1}{\sqrt{5}} \begin{bmatrix} -1 \\ 2 \\ 0 \end{bmatrix}$  is an orthonormal basis for  $W^\perp$ .

**Alternatively.** If we didn't see that, we can use that  $W = \text{col} \left( \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \right)$  so that  $W^\perp = \text{null} \left( \begin{bmatrix} 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right)$ .

The latter has basis  $\begin{bmatrix} -1/2 \\ 1 \\ 0 \end{bmatrix}$ , which becomes  $\frac{1}{\sqrt{5}} \begin{bmatrix} -1 \\ 2 \\ 0 \end{bmatrix}$  when normalized.

In conclusion, we have  $A = PDP^T$  with  $P = \begin{bmatrix} 2/\sqrt{5} & 0 & -1/\sqrt{5} \\ 1/\sqrt{5} & 0 & 2/\sqrt{5} \\ 0 & 1 & 0 \end{bmatrix}$  and  $D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$ .

(b) Here,  $W = \text{span} \left\{ \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$ .

- (basis for  $W$ ) We apply Gram-Schmidt:

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - \frac{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}} \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} - \frac{3}{5} \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} -1 \\ 5 \\ 2 \end{bmatrix}$$

Normalizing, we find that  $\frac{1}{\sqrt{5}} \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \frac{1}{\sqrt{30}} \begin{bmatrix} -1 \\ 5 \\ 2 \end{bmatrix}$  is an orthonormal basis for  $W$ .

- (basis for  $W^\perp$ ) Since  $W = \text{col} \left( \begin{bmatrix} 2 & 1 \\ 0 & 1 \\ 1 & 1 \end{bmatrix} \right)$ , we have  $W^\perp = \text{null} \left( \begin{bmatrix} 2 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix} \right)$ .

Solving the system,  $\begin{bmatrix} 2 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix} \rightsquigarrow \begin{bmatrix} 2 & 0 & 1 \\ 0 & 1 & 1/2 \end{bmatrix}$ , we find that  $\begin{bmatrix} -1/2 \\ -1/2 \\ 1 \end{bmatrix}$  is a basis for  $W^\perp$ . Normalized:  $\frac{1}{\sqrt{6}} \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$

**Alternatively.** We could also use  $W = \text{col} \left( \begin{bmatrix} 2 & -1 \\ 0 & 5 \\ 1 & 2 \end{bmatrix} \right)$  so that  $W^\perp = \text{null} \left( \begin{bmatrix} 2 & 0 & 1 \\ -1 & 5 & 2 \end{bmatrix} \right)$ .

In conclusion, we have  $A = PDP^T$  with  $P = \begin{bmatrix} 2/\sqrt{5} & -1/\sqrt{30} & -1/\sqrt{6} \\ 0 & 5/\sqrt{30} & -1/\sqrt{6} \\ 1/\sqrt{5} & 2/\sqrt{30} & 2/\sqrt{6} \end{bmatrix}$  and  $D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$ .

### Problem 8.

- (a) The eigenvalues of a  $5 \times 5$  matrix for orthogonally projecting onto a 3-dimensional subspace are .

What are the eigenspaces of that matrix?

- (b) Suppose  $A$  is the  $3 \times 3$  matrix of a reflection through a plane (containing the origin).

